

LARGE CARDINALS AND GAP-1 MORASSES

ANDREW D. BROOKE-TAYLOR AND SY-DAVID FRIEDMAN

Abstract. We present a new partial order for directly forcing morasses to exist that enjoys a significant homogeneity property. We then use this forcing in a reverse Easton iteration to obtain an extension universe with morasses at every regular uncountable cardinal, while preserving all n -superstrong ($1 \leq n \leq \omega$), hyperstrong and 1-extendible cardinals. In the latter case, a preliminary forcing to make the GCH hold is required. Our forcing yields morasses that satisfy an extra property related to the homogeneity of the partial order; we refer to them as *mangroves* and prove that their existence is equivalent to the existence of morasses. Finally, we exhibit a partial order that forces universal morasses to exist at every regular uncountable cardinal, and use this to show that universal morasses are consistent with n -superstrong, hyperstrong, and 1-extendible cardinals. This all contributes to the second author's *outer model programme*, the aim of which is to show that L -like principles can hold in outer models which nevertheless contain large cardinals.

§1. Introduction. Morasses are combinatorial structures formulated by Jensen to abstract out properties of L useful for proving cardinal transfer theorems. Originally thought to be such complex objects that one should not attempt to understand them outside the context of their construction in L (according to [11]), they were opened up to broader study and deeper understanding by Velleman [12], who amongst other things presented a particularly elegant partial order with which one can force morasses to exist. This lends itself to use in the second author's *outer model programme* in which, as a counterpoint to the well-known inner model programme, the goal is to force to obtain L -like outer models for various large cardinals. Indeed, in [7] it is shown that one may force morasses to exist at every regular uncountable cardinal, while preserving a given n -superstrong or hyperstrong cardinal.

In this article we extend these results. In addition to n -superstrong and hyperstrong cardinals we consider 1-extendible cardinals, for which a careful consideration of forcing the GCH as a preliminary step is appropriate. Further, we exhibit a new forcing partial order to obtain the morasses, which allows us to preserve *all* cardinals of the considered kinds. This is achieved by modifying Velleman's forcing to give it a degree of homogeneity. Finally, we show how one may force universal morasses to exist. This new forcing again

Andrew Brooke-Taylor was supported for this research by Austrian Science Fund (FWF) project P16790-N04.

Sy-David Friedman was supported by FWF projects P16790-N04 and P19375-N18.

fits in with the techniques of [7], and so we obtain that universal morasses can be forced while preserving an n -superstrong, hyperstrong or 1-extendible cardinal. This answers a question of Pereira to the authors.

The reader may be aware that the existence of morasses has been shown by Velleman [13] to be equivalent to the existence of much simpler objects called *simplified morasses*. However, while these latter structures are much more manageable, the only known way to obtain them in L is via morasses themselves. Thus, to stay more directly in keeping with the goal of forcing L -like models, we stick with morasses. On the other hand, we do use Velleman's [12] axiomatisation of morasses, which already strips away some of the detail of morasses as defined by Jensen. This will have ramifications when we force to obtain universal morasses — see section 9 below.

§2. Preliminaries: the GCH and 1-Extendible Cardinals.

Our forcing constructions will rely on having the GCH hold in the ground model, so we discuss here how to attain it by a preliminary forcing. The iterated forcing we use to obtain the GCH is a fairly standard, natural one, at each stage collapsing the old 2^{\aleph_α} to $\aleph_{\alpha+1}$. The main issue is to show that forcing with this particular partial order does preserve the large cardinals we are interested in. In many cases, this was achieved in [7]; we show here that similar arguments go through for 1-extendible cardinals.

Following [10], we denote by $\text{Fn}(I, J, \lambda)$ the set of partial functions from I to J with cardinality less than λ .

DEFINITION 1. *The GCH Partial Order P is the reverse Easton iteration of $\langle \dot{Q}_\alpha \mid \alpha \in \text{Ord} \rangle$, in which direct limits are taken only at strongly inaccessible cardinals and inverse limits are taken at other limit stages, where \dot{Q}_α is the canonical P_α -name for $\text{Fn}(\aleph_{\alpha+1}, 2, \aleph_{\alpha+1})$, with $\aleph_{\alpha+1}$ to be evaluated in $V[G_\alpha]$.*

For the rest of this section $P = P_{\text{Ord}}$ will denote the GCH Partial Order, with P_α denoting the iteration after α stages, $G = G_{\text{Ord}}$ denoting a P -generic over V , G_α denoting $G \restriction P_\alpha$, and Q_α denoting $\text{Fn}(\aleph_{\alpha+1}, 2, \aleph_{\alpha+1})$ in $V[G_\alpha]$.

Before proceeding with the analysis of this iteration, let us recall the basic properties of the forcing poset $\text{Fn}(\aleph_{\alpha+1}, 2, \aleph_{\alpha+1})$. It is $\aleph_{\alpha+1}$ -closed and $(2^{\aleph_\alpha})^+$ -cc, so the only cardinals collapsed by forcing with it are those κ such that $\aleph_{\alpha+1} < \kappa \leq 2^{\aleph_\alpha}$. Of course, since $\text{Fn}(\aleph_{\alpha+1}, 2, \aleph_{\alpha+1})$ is equivalent to $\text{Fn}(\aleph_{\alpha+1}, 2^{\aleph_\alpha}, \aleph_{\alpha+1})$, a surjection $\aleph_{\alpha+1} \rightarrow 2^{\aleph_\alpha}$ is added, and all such κ are indeed collapsed. By a nice names argument, the continuum function is unchanged by this forcing at and above the ground model 2^{\aleph_α} , and $2^{\aleph_{\alpha+1}}$ in the extension will be the ground model $2^{2^{\aleph_\alpha}}$. By $\aleph_{\alpha+1}$ -closure, the continuum function is unchanged below \aleph_α , and after the forcing we have $2^{\aleph_\alpha} = \aleph_{\alpha+1}$.

Now, to the GCH partial order itself. To demonstrate that ZFC is preserved by this class forcing, we show that it is tame (see [6]).

LEMMA 2. *The GCH Partial Order is tame.*

PROOF. By the Factor Lemma (see for example [8], Lemma 21.8), P may be written as $P_\alpha * \dot{P}^\alpha$, where \dot{P}^α names the iteration starting with Q_α . Each iterand in \dot{P}^α is forced to be $\aleph_{\alpha+1}$ -closed, and inverse limits are taken everywhere except at inaccessibles of V , which will remain inaccessible after forcing with P_α (indeed, it can be shown by induction that P_α has a dense suborder of size at most $\beth_{\alpha+1}^V$).

We therefore have that for all α

$$1_{P_\alpha} \Vdash \dot{P}^\alpha \text{ is } \aleph_{\alpha+1}\text{-closed},$$

and this implies that P is tame — see [6], Lemmas 2.22 and 2.31. \dashv

LEMMA 3. *Let α be an ordinal or Ord. Then for every ordinal $\gamma \leq \alpha$, $\beth_\gamma^{V[G_\alpha]} = \aleph_\gamma^{V[G_\alpha]}$.*

PROOF. As in the proof of Lemma 2 above, P_α may be factorised as $P_\gamma * \dot{P}^{[\gamma, \alpha]}$, where $\dot{P}^{[\gamma, \alpha]}$ names the iteration starting with Q_γ , and we have

$$1_{P_\gamma} \Vdash \dot{P}^{[\gamma, \alpha]} \text{ is } \aleph_{\gamma+1}\text{-closed}.$$

It hence suffices to show that $\beth_\gamma^{V[G_\gamma]} = \aleph_\gamma^{V[G_\gamma]}$ for every γ . This may be proven by induction on γ , with the successor step following from the discussion of $\text{Fn}(\aleph_{\alpha+1}, 2, \aleph_{\alpha+1})$ above. \dashv

In particular, these two lemmas combine to show that forcing with P yields a model of ZFC + GCH.

In [7], it is shown that when forcing with the GCH forcing P , any given hyperstrong or n -superstrong cardinal may be preserved. In fact, the same proof essentially gives the following stronger statement.

THEOREM 4. *Forcing with the GCH partial order P preserves all hyperstrong cardinals and n -superstrong cardinals, for all $n \in \omega + 1$.*

PROOF. Observe that each iterand in the GCH forcing is very homogeneous: given a condition $p \in Q_\alpha = \text{Fn}(\aleph_{\alpha+1}, 2, \aleph_{\alpha+1})$ and a Q_α -generic g , there is an automorphism π of Q_α such that $p \in \pi''g$. Such strong homogeneity is known to hold for an entire iteration if it is forced to hold for each iterand; see [5] for the details. It follows that all the arguments for Theorem 4.2 of [7], which in particular choose P -generics containing a certain master condition for each large cardinal κ , can be carried out within $V[G]$. Therefore, all large cardinals of the listed kinds are preserved in $V[G]$. \dashv

We shall show that the same result is true for 1-extendible cardinals. Recall their definition:

DEFINITION 5. *A cardinal κ is 1-extendible if there is a $\lambda > \kappa$ and an elementary embedding $j : H_{\kappa^+} \rightarrow H_{\lambda^+}$ with critical point κ .*

Equivalently, κ is 1-extendible if and only if there is a $\lambda > \kappa$ and an elementary embedding $V_{\kappa+1} \rightarrow V_{\lambda+1}$ with critical point κ . The above definition will be more convenient for our purposes, however, as $H_{\kappa+}$ forms a model of ZFC minus the Power Set axiom, which we shall refer to as ZFC^- . 1-extendible cardinals lie in consistency strength between superstrong and hyperstrong cardinals (see [9], Proposition 26.11) and play an interesting role in the outer model programme, as Cummings and Schimmerling [3] have shown them to be essentially the strongest large cardinals compatible with the principle \square .

THEOREM 6. *For any model V of ZFC , there is a class-generic extension $V[G]$ of V such that $V[G] \models GCH$, and in which every 1-extendible cardinal of V remains 1-extendible.*

PROOF. We shall of course force with the GCH forcing P . Let κ be 1-extendible in V , let $j : H_{\kappa+} \rightarrow H_{\lambda+}$ in V witness this fact, and let G be P -generic over V . We shall show that j may be lifted to an embedding $j^* : H_{\kappa+}^{V[G]} \rightarrow H_{\lambda+}^{V[G]}$ witnessing the 1-extendibility of κ in $V[G]$.

The GCH forcing P may be factorised as $P_{\kappa} * P^{\kappa}$, with P^{κ} a κ^+ -closed forcing. Hence, $H_{\kappa+}^{V[G]} = H_{\kappa+}^{V[G_{\kappa}]}$. Similarly, $H_{\lambda+}^{V[G]} = H_{\lambda+}^{V[G_{\lambda}]}$. Now because κ is inaccessible, P_{κ} has cardinality κ and lies in $H_{\kappa+}$. Recall that forcing over a model of ZFC^- yields a model of ZFC^- , and note that G_{κ} is generic over $H_{\kappa+}$ for the set forcing P_{κ} . We claim that in fact $H_{\kappa+}^{V[G_{\kappa}]} = H_{\kappa+}^V[G_{\kappa}]$, the generic extension of the model $H_{\kappa+}^V$ of ZFC^- by G_{κ} . For this we need to show that if σ is a P_{κ} -name in V such that $\sigma_{G_{\kappa}} \in H_{\kappa+}^{V[G_{\kappa}]}$, then there is a name $\tau \in H_{\kappa+}^{P_{\kappa}}$ such that $\sigma_{G_{\kappa}} = \tau_{G_{\kappa}}$. But now every element of $H_{\kappa+}$ can be obtained as the Mostowski collapse of a relation on κ given by a subset of $\kappa \times \kappa$ (and this process does not appeal to the Power Set Axiom). Every subset of $\kappa \times \kappa$ in the extension $V[G_{\kappa}]$ has a nice name of the form

$$\bigcup_{(\alpha, \beta) \in \kappa \times \kappa} \{(\check{\alpha}, \check{\beta})\} \times A_{(\alpha, \beta)}$$

where each $A_{(\alpha, \beta)}$ is an antichain in P_{κ} . Since $|P_{\kappa}| = \kappa$, such nice names lie in $H_{\kappa+}^V$. Therefore, every subset of $\kappa \times \kappa$ in $V[G_{\kappa}]$ is also in $H_{\kappa+}^V[G_{\kappa}]$, and consequently we indeed have that $H_{\kappa+}^{V[G_{\kappa}]} = H_{\kappa+}^V[G_{\kappa}]$. Of course by elementarity, λ is also inaccessible, and so also $H_{\lambda+}^{V[G_{\lambda}]} = H_{\lambda+}^V[G_{\lambda}]$.

We are thus reduced to showing that $j : H_{\kappa+} \rightarrow H_{\lambda+}$ may be lifted to an elementary embedding $j^* : H_{\kappa+}^V[G_{\kappa}] \rightarrow H_{\lambda+}^V[G_{\lambda}]$. We wish to apply Silver's technique of lifting embeddings (see [2], Section 9), and must show that $j''G_{\kappa} \subseteq G_{\lambda}$. But now $j \restriction P_{\kappa}$ is the identity function, so $j''G_{\kappa} = G_{\kappa} \subset G_{\lambda}$. Therefore j indeed lifts to $j^* : H_{\kappa+}^{V[G]} \rightarrow H_{\lambda+}^{V[G]}$, and so κ is 1-extendible in $V[G]$. \dashv

In Section 8, we will show that 1-extendible cardinals κ may be preserved while forcing morasses to exist. To show that the forcing

is pretame from the perspective of $H_{\kappa+}$, we will first force GCH as above, and use the fact that after our GCH forcing, $V[G] = L[G]$, and further $H_{\kappa+}^{V[G]} = L_{\kappa+}[A]$, where A is taken to be a class predicate over $H_{\kappa+}$ for the Cohen set added at stage κ . This gives a stratification of our model $H_{\kappa+}$ of ZFC^- into sets, as is required for the proof that the forcing relation is definable for a pretame class forcing — see [6], Theorem 2.18. Only then will we be able to conclude that the forcing relation for our forcing is definable.

However, $p \Vdash \varphi(\sigma_0, \dots, \sigma_{n-1})$ will only be definable relative to A , and so the usual lifting lemma will not suffice to lift the embedding j witnessing 1-extendibility, as j need not respect arbitrary class predicates. We show here that in fact j *does* respect the predicate A given by the generic, giving a mild strengthening of Theorem 6. The crux of the proof will be the definability of the forcing relation for $P_{\kappa+1}$, which from the point of view of $H_{\kappa+}$ is a class forcing. The usual argument for pretame class forcing (Theorem 2.18 of [6]) is not applicable, since we don't yet have a stratification of $H_{\kappa+}$ into sets. Nevertheless, we will be able to demonstrate the definability of forcing for $P_{\kappa+1}$ because it is such a well-behaved forcing partial order.

DEFINITION 7. *The language \mathcal{L}_{STG} is the language obtained from the language of set theory by adding a single unary predicate G .*

LEMMA 8. *Let $\kappa \in M$ be an inaccessible cardinal, and let Q_κ denote $\text{Fn}(\kappa^+, 2, \kappa^+)$, the forcing at stage κ in the GCH forcing above. Then the forcing relation $\Vdash \varphi$ for forcing over $H_{\kappa+}$ with Q_κ is definable over $H_{\kappa+}$ for \mathcal{L}_{STG} formulae φ , where the predicate G is to be interpreted as the Q_κ -generic, and the Truth Lemma holds: a formula is true in the extension if and only if it is forced by some condition in the generic.*

PROOF. We follow the expository style of [10], defining a relation $\Vdash^* \varphi$ for each φ , which will then be seen to be equivalent to $\Vdash \varphi$. We will make use of the closure properties of Q_κ to make this definition for atomic formulae, from which point it may be extended to all formulae as usual. As a first step, we give a definition for forcing the predicate G to hold: define $p \Vdash^* G(\sigma)$ to mean

$$(1) \quad \forall q \leq p \exists r \leq q \exists s \geq r (r \Vdash^* \sigma = \check{s}),$$

that is, it is dense below p to force σ to be something specific greater than yourself. If $p \in G$, then by genericity there is some $r \in G$ and some s greater than or equal to r and consequently also in G such that $r \Vdash^* \sigma = \check{s}$, whence $\sigma_G \in G$. Conversely, suppose $\sigma_G \in G$ for every $G \ni p$; let $q \leq p$ and fix some $G \ni q$. Once we have shown that the forcing relation is definable for atomic formulae, it will follow from the truth lemma there is some $r' \in G$ such that

$$r' \Vdash \sigma = (\check{\sigma}_G).$$

Taking $r \in G$ such that $r \leq q$, $r \leq r'$, and $r \leq \sigma_G$, we have that r makes (1) hold with σ_G as s . Hence, (1) is indeed a formal definition encapsulating the statement that for every generic containing p , σ_G is in G , and it is clear that the argument extends to prove the Truth Lemma valid for formulae involving the predicate G .

With forcing the predicate G so defined, we can move on to show that the forcing relation is indeed definable for atomic formulae. Let σ and τ be Q_κ -names in H_{κ^+} , and let

$$R_{\sigma,\tau} = \text{trcl}(\{\sigma, \tau\}) \cap P,$$

the set of conditions hereditarily appearing in either σ or τ . Note that since the names σ and τ are in H_{κ^+} , $|R_{\sigma,\tau}| \leq \kappa$. Let $d_{\sigma,\tau} = \bigcup_{r \in R_{\sigma,\tau}} \text{dom}(r)$; then $d_{\sigma,\tau}$ is a subset of κ^+ of size at most κ .

Now, for conditions q such that $d_{\sigma,\sigma} \subseteq \text{dom}(q)$, we may recursively define the evaluation of σ at q by

$$\text{val}(\sigma, q) = \{\text{val}(\rho, q) \mid \exists X \subseteq \text{dom}(q) (\langle \rho, q \restriction X \rangle \in \sigma)\}.$$

Observe that if G is any Q_κ -generic over H_{κ^+} containing such a q , then $\sigma_G = \text{val}(\sigma, q)$. Likewise, for any G which is Q_κ -generic over H_{κ^+} and any element q of G with domain containing $d_{\sigma,\sigma}$, we have $\text{val}(\sigma, q) = \sigma_G$. Thus, the evaluation of any name in the generic extension may be entirely determined by a single condition.

With this in hand, we may define

$$p \Vdash^* \sigma \in \tau \iff \forall q \leq p (\text{dom}(q) \supseteq d_{\sigma,\tau} \rightarrow \text{val}(\sigma, q) \in \text{val}(\tau, q))$$

and

$$p \Vdash^* \sigma = \tau \iff \forall q \leq p (\text{dom}(q) \supseteq d_{\sigma,\tau} \rightarrow \text{val}(\sigma, q) = \text{val}(\tau, q)).$$

Then clearly $p \Vdash^* \sigma \in \tau$ if and only if $\sigma_G \in \tau_G$ for every Q_κ -generic $G \ni p$, and $p \Vdash^* \sigma = \tau$ if and only if $\sigma_G = \tau_G$ for every Q_κ -generic $G \ni p$. Moreover, if $\sigma_G \in \tau_G$, then any $p \in G$ with $\text{dom}(p) \supseteq d_{\sigma,\tau}$ will force $\sigma \in \tau$, and the Truth Lemma follows. \dashv

We are now ready for our strengthening of Theorem 6. Let us denote by $G(\kappa)$ the generic from stage κ of G , that is, the Q_κ -generic associated to G .

THEOREM 9. *Let V be a model of ZFC and let κ be a 1-extendible cardinal in V with 1-extendibility witnessed by an elementary embedding $j : H_{\kappa^+} \rightarrow H_{\lambda^+}$. Let $V[G]$ be a P -generic extension of V . Then there is a $G' \subset V[G]$ which is P -generic over V such that $V[G] = V[G']$ and the lift j^* of j to $H_{\kappa^+}^{V[G']}$ (as in the proof of Theorem 6) is elementary between the \mathcal{L}_{STG} -structures $\langle H_{\kappa^+}^{V[G]}, G'(\kappa) \rangle$ and $\langle H_{\lambda^+}^{V[G]}, G'(\lambda) \rangle$.*

PROOF. In Theorem 6 the lift j^* was constructed in $H_{\kappa^+}^{V[G_\kappa]}$, after observing that P^κ is κ^+ -closed and hence does not affect H_{κ^+} . We now claim that the P^κ generic G^κ may be chosen so that this same j^* is also elementary for formulae in the language \mathcal{L}_{STG} . By the

Truth Lemma, every \mathcal{L}_{STG} sentence φ true in $\langle H_{\kappa^+}^{V[G_{\kappa+1}]}, G(\kappa) \rangle = \langle H_{\kappa^+}^{V[G]}, G(\kappa) \rangle$ is forced (over $H_{\kappa^+}^{V[G_\kappa]}$) to be true by some $p \in Q_\kappa$. Now by the definability of the forcing relation, we have

$$p \Vdash_{Q_\kappa} \varphi(\sigma_0, \dots, \sigma_n) \leftrightarrow j^*(p) \Vdash_{Q_\lambda} \varphi(j^*(\sigma_0), \dots, j^*(\sigma_n)).$$

Therefore, if $j^* "G(\kappa) \subseteq G(\lambda)$, we will have that j^* is elementary from $\langle H_{\kappa^+}^{V[G]}, G(\kappa) \rangle$ to $\langle H_{\lambda^+}^{V[G]}, G(\lambda) \rangle$. But now $|G(\kappa)|^{V[G_\lambda]} = (\kappa^+)^{V[G_\lambda]} < \lambda$, so $j^* "G(\kappa)$ is a pairwise-compatible set of conditions in $\text{Fn}(\lambda, 2, \lambda)^{V[G_\lambda]}$ (indeed, in $\text{Fn}(\lambda, 2, \lambda)^{V[G_\kappa]}$) of size less than λ . Moreover, this set lies in $V[G_\lambda]$, since j , G_κ and $G(\kappa)$ all do. Hence, $m = \bigcup j^* "G(\kappa)$ is a single condition in $\text{Fn}(\lambda, 2, \lambda)^{V[G_\lambda]} = Q_\lambda$, and so if G is chosen such that this master condition lies in $G(\lambda)$, then we will indeed have $j^* "G(\kappa) \subset G(\lambda)$. This is easy to arrange by modifying $G(\lambda)$ to obtain $G'(\lambda)$. Let q be the element of $G(\lambda)$ with $\text{dom}(q) = \text{dom}(m)$, and define $\psi : Q_\lambda \rightarrow Q_\lambda$ changing partial functions by switching their values between 0 and 1 on points where q and m disagree, that is,

$$\psi(f)(\alpha) = \begin{cases} 0 & \text{if } q(\alpha) \neq m(\alpha) \text{ and } f(\alpha) = 1 \\ 1 & \text{if } q(\alpha) \neq m(\alpha) \text{ and } f(\alpha) = 0 \\ f(\alpha) & \text{otherwise} \end{cases}$$

for all $f \in Q_\lambda$ and $\alpha \in \text{dom}(f)$. Clearly ψ is an (involutive) automorphism of Q_λ in $V[G]$, whence $G'(\lambda) = \psi "G(\lambda)$ is Q_λ -generic over $V[G_\lambda]$, and $V[G_{\lambda+1}] = V[G_\lambda * G'(\lambda)]$. The “tail” generic $G^{\lambda+1}$ will of course still be $P^{\lambda+1}$ -generic over $V[G_\lambda * G'(\lambda)]$. Hence, considering P as $P_\lambda * Q_\lambda * P^{\lambda+1}$, we may take $G' = G_\lambda * G'(\lambda) * G^{\lambda+1}$, which is P -generic over V and satisfies $V[G] = V[G']$. Moreover, $G'(\lambda)$ contains the master condition m determined by $G(\kappa) = G'(\kappa)$, so $j^* "G'(\kappa) \subset G'(\lambda)$, and so j^* is elementary from $\langle H_{\kappa^+}^{V[G']}, G'(\kappa) \rangle$ to $\langle H_{\lambda^+}^{V[G']}, G'(\lambda) \rangle$, as required. \dashv

§3. Basic definitions. The definition for morasses that we use will be that of Velleman [12]. In particular, we retain the notation M.1–M.7 for axioms of a morass, used both there and in Devlin [4]. We also follow Velleman in separating out the following subsidiary definition, although we change the terminology slightly to emphasise the order preservation property.

DEFINITION 10. *A function $\pi : \alpha \rightarrow \beta$ between two ordinals is a successor, limit, zero and order preserving (SLOOP) function if*

- for all $\gamma < \alpha$, $\pi(\gamma + 1) = \pi(\gamma) + 1$, and
- for all limit $\gamma < \alpha$, $\pi(\gamma)$ is also a limit, and
- $\pi(0) = 0$, and
- for all $\gamma < \delta < \alpha$, $\pi(\gamma) < \pi(\delta)$.

Note in particular that the composition of SLOOP functions will yield a SLOOP function.

Morasses will be defined based on a set \mathcal{S} of ordered pairs of ordinals. For such an ordered pair $x = \langle \alpha, \beta \rangle$, let us denote α by $l(x)$ (the *level* of x), and β by $o(x)$ (the *order* of x). Also, note that we here use the word “tree” in the liberal sense where others might use “forest”: our tree will have many root nodes.

DEFINITION 11. *For any uncountable regular cardinal κ , a $(\kappa, 1)$ -morass (or simply morass when κ is clear from the context) consists of:*

- i. *a subset \mathcal{S} of $(\kappa \times \kappa) \cup (\{\kappa\} \times \kappa^+)$, and*
- ii. *A tree order \prec on \mathcal{S} , and*
- iii. *For every pair $\langle x, y \rangle$ of elements of \mathcal{S} with $x \prec y$, a function $\pi_{xy} : o(x) + 1 \rightarrow o(y) + 1$,*

such that the following conditions hold.

Left-alignment: *For each $\alpha \leq \kappa$, let $\theta_\alpha = \{\beta \mid \langle \alpha, \beta \rangle \in \mathcal{S}\}$. Then θ_α is in fact an ordinal. Moreover, $\theta_\kappa = \kappa^+$, and for $\alpha < \kappa$, $0 < \theta_\alpha < \kappa$.*

Monotonicity: *For x and y in \mathcal{S} , $x \prec y$ implies $l(x) < l(y)$.*

Commutativity: *If $x \prec y \prec z$ then $\pi_{yz} \circ \pi_{xy} = \pi_{xz}$.*

M.1: *For each pair $x, y \in \mathcal{S}$ with $x \prec y$, π_{xy} is an SLOOP function, and $\pi_{xy}(o(x)) = o(y)$.*

M.2: *Suppose $x \prec y \in \mathcal{S}$ and $\nu < o(x)$. Let $w = \langle l(x), \nu \rangle$ and $z = \langle l(y), \pi_{xy}(\nu) \rangle$. Then $w \prec z$ and $\pi_{wz} = \pi_{xy} \upharpoonright (\nu + 1)$.*

M.3: *For all $y \in \mathcal{S}$, $\{l(x) \mid x \prec y\}$ is closed in $l(y)$.*

M.4: *For all $y \in \mathcal{S}$, if $o(y) + 1 \neq \theta_{l(y)}$ (as defined for Left-alignment above), then $\{l(x) \mid x \prec y\}$ is unbounded in $l(y)$. In particular, if α is a successor ordinal, then $\theta_\alpha = 1$.*

M.5: *For all $y \in \mathcal{S}$, if $\{l(x) \mid x \prec y\}$ is unbounded in $l(y)$, then $o(y) = \bigcup \{\pi_{xy}“o(x) \mid x \prec y\}$.*

M.6: *Suppose $x \prec y \in \mathcal{S}$ and $o(x)$ is a limit ordinal. Let $\nu = \sup(\pi_{xy}“o(x))$ and let $z = \langle l(y), \nu \rangle$. Then $x \prec z$ and $\pi_{xz} \upharpoonright o(x) = \pi_{xy} \upharpoonright o(x)$.*

M.7: *Suppose $x \prec y \in \mathcal{S}$, $l(x) < \alpha < l(y)$, $o(x)$ is a limit ordinal, and $o(y) = \sup(\pi_{xy}“o(x))$. If*

$$\forall \nu < o(x) \exists \gamma (\langle \alpha, \gamma \rangle \prec \langle l(y), \pi_{xy}(\nu) \rangle),$$

then there is a γ such that $\langle \alpha, \gamma \rangle \prec y$.

Note in particular axiom M.5, in the case where $l(y) = \kappa$: for any $\tau < \kappa^+$, we have τ expressed as the (increasing, by commutativity) union of the images of the maps $\pi_{x\langle \kappa, \tau \rangle} \upharpoonright o(x)$ for $x \prec \langle \kappa, \tau \rangle$. Further note that this isn’t just a variant of κ many things “adding up” to κ , and then τ being bijective with κ , but something more direct: the maps $\pi_{x\langle \kappa, \tau \rangle}$ are order preserving, and so the ordinals mapping into τ must to some extent reflect the structure of τ . For example, if τ is a successor ordinal and so has a largest element, then for $x \prec \langle \kappa, \tau \rangle$ with $l(x)$ sufficiently large, $o(x)$ must also have a largest element. Jensen’s original definition in fact called for even stronger

preservation properties; see [4] for details. Also observe that for any $x \preceq y$ in \mathcal{S} , $o(x) \leq o(y)$ since by M.1, π_{xy} is a strictly order-preserving function from $o(x) + 1$ to $o(y) + 1$.

When we force morasses to exist in the presence of multiple large cardinals, it will be convenient for the sake of preservation of the large cardinal property to use a partial order which lends itself to homogeneity arguments (see Section 6 below). The upshot will be that the generic morass satisfies a useful property not possessed by the morasses obtained by forcing with Velleman's partial order defined in [12]. We give here a name for the kind of morass that we obtain.

DEFINITION 12. *Suppose $M = \langle \mathcal{S}, \preceq, \langle \pi_{xy} \rangle_{x \preceq y} \rangle$ is a $(\kappa, 1)$ -morass. An ordinal $\alpha < \kappa$ is a mangal of M if for all $x, y \in \mathcal{S}$ with $x \preceq y$ and $l(x) < \alpha < l(y)$, there is a $z \in \mathcal{S}$ with $l(z) = \alpha$ such that $x \preceq z \preceq y$. If the set of mangals of M is cofinal in κ , we say that M is a κ -mangrove, or simply mangrove when κ is clear from the context.*

An immediate question is whether the existence of a morass implies the existence of a mangrove. It turns out that the answer is yes, as we shall show in Section 5, using Velleman's theorem that the existence of a morass is equivalent to a certain forcing axiom.

Another natural question is what the set of mangals of a morass can look like. A first result in this direction is the following.

PROPOSITION 13. *For any morass M , the set of mangals of M is closed in κ .*

PROOF. Suppose that $\gamma < \kappa$ is a limit point of the set of mangals of a morass $M = \langle \mathcal{S}, \preceq, \pi \rangle$, and let $x, y \in \mathcal{S}$ satisfy $x \preceq y$ and $l(x) < \gamma < l(y)$. For each mangal α of M such that $l(x) < \alpha < \gamma$, there is a z_α with $l(z_\alpha) = \alpha$ and $x \preceq z_\alpha \preceq y$. But then by M.3, there must be some z_γ with $l(z_\gamma) = \gamma$ and $x \preceq z_\gamma \preceq y$; thus, γ is also a mangal of M . \dashv

In particular, if M is a mangrove, the set of mangals of M is a closed unbounded subset of κ .

§4. Forcing morasses to exist. As is the case for many combinatorial structures, one can force with a partial order of partial morasses to get a morass in the generic extension, as we shall show below. We use a partial order similar to those described in [7] and [12]. However, ours will differ in that we strengthen the requirements for an extension of a condition, ensuring that each condition of the generic goes up to a mangal of the ultimate morass. Naturally, this will make our generic morass a mangrove, whereas it is not hard to check that a generic morass for Velleman's partial order in [12] will be far from a mangrove; indeed, it will be κ -branching at the node $\langle 0, 0 \rangle$, and so cannot have any mangals at all! However, many of the

details of our proof will remain essentially the same as in that paper. In particular, we repeatedly use the basic construction given there for extending morass conditions, tweaking it to fit the particular requirements in each case.

4.1. Definitions. Let $x \prec_i y$ denote that y is an immediate \prec -successor of x .

DEFINITION 14. *For any uncountable regular cardinal κ , a $(\kappa, 1)$ -morass condition consists of:*

- i. *a subset \mathcal{S} of $((\lambda + 1) \times \kappa) \cup (\{\kappa\} \times \kappa^+)$ for some $\lambda < \kappa$, and*
- ii. *A tree order \prec on \mathcal{S} , and*
- iii. *For every pair $\langle x, y \rangle$ of elements of \mathcal{S} with $x \prec y$, a function $\pi_{xy} : o(x) + 1 \rightarrow o(y) + 1$,*

such that

- 1. *Left-alignment holds for all $\alpha \leq \lambda$, and the set $S = \{\beta \mid \langle \kappa, \beta \rangle \in \mathcal{S}\}$ contains 0 and is closed under ordinal successors and predecessors.*
- 2. *Let f be the order-preserving bijection from $ot(S)$ to S . Then $ot(S) \leq \theta_\lambda$, and for each $\nu < ot(S)$, $\langle \lambda, \nu \rangle \prec \langle \kappa, f(\nu) \rangle$.*
- 3. *Monotonicity, Commutativity, M.1, M.2, M.3 and M.5 hold. Axiom M.4 holds for those $y \in \mathcal{S}$ such that $l(y) \leq \lambda$. Axioms M.6 and M.7 hold for those x and y such that $x \prec_i y$ and $l(y) \leq \lambda$.*

For our analysis, we will need to extend the new notion of a mangal to also be applicable to morass conditions.

DEFINITION 15. *Let $p = \langle \mathcal{S}, \prec, \pi \rangle$ be a morass condition. An ordinal $\alpha \leq \lambda$ is a mangal of p if for all $x, y \in \mathcal{S}$ with $x \prec y$ and $l(x) < \alpha < l(y)$, there is a $z \in \mathcal{S}$ with $l(z) = \alpha$ such that $x \prec z \prec y$.*

Our definition of a morass condition is the same as that used in [12]; our partial order will differ in the definition of \leq . For this reason, we continue to refer to these conditions as *morass* conditions, even though we shall call our partial order the *mangrove* forcing. Also note that, since it arises frequently and the meaning is fairly clear, we shall consistently abuse notation, writing $x \in p$ to mean that x is an element of the set \mathcal{S} for the morass condition p .

A few comments about the definition of morass conditions are in order at this point. Note that requiring Axiom M.2 when $l(y) = \kappa$ implicitly entails imposing the condition on S and the various π_{xy} that for any $x \prec y$ with $l(y) = \kappa$, $\pi_{xy} " (o(x) + 1) \subset S$. Requiring M.7 for $x \prec_i y$ is the same as positing the non-existence of α fitting the antecedent of that axiom, since $x \prec \langle \alpha, \gamma \rangle \prec y$ contradicts $x \prec_i y$.

Observe that by requirement 1 above, $ot(S)$ must be a limit ordinal; hence by requirement 2, $\theta_\lambda > 1$, and so by M.4, λ is a limit ordinal. Also, from M.2, requirement 2, and the fact that \prec is a tree order, we get that for $x \prec y$ with $l(x) = \lambda$ and $l(y) = \kappa$, $\pi_{xy} = f \upharpoonright o(x) + 1$.

Similarly, because \prec is required to be a tree order and f is surjective, one can only have $x \prec y$ with $l(y) = \kappa$ when either $l(x) = \lambda$, or there is a z such that $x \prec z \prec y$ with $l(z) = \lambda$. Thus, λ is a mangal of the morass condition; this basic example of a mangal will be important in later arguments. As a result of λ being a mangal, it is equivalent to only explicitly require M.3 to hold for those $y \in \mathcal{S}$ with $l(y) < \kappa$. The same is trivially true of M.5, which for morass conditions is vacuous in the $l(y) = \kappa$ case. Finally, note that because of the closure properties of S given in requirement 1, f and f^{-1} are SLOOP functions.

For any regular cardinal κ , let P_κ be the set of $(\kappa, 1)$ -morass conditions, along with an extra point $\mathbf{1}$ to act as a maximum element; we will generally ignore $\mathbf{1}$, it being trivial to extend the definitions to encompass it as a special case. For each non- $\mathbf{1}$ element p of P_κ , let us denote with superscript p the components and defined notions of p : \mathcal{S}^p , \prec^p , π_{xy}^p for $x \prec^p y$, λ^p , S^p as in requirement 1, f^p as in requirement 2, and θ_α^p . We shall also refer to $\{x \in p \mid l(x) = \lambda^p\}$ as the *top level* of the condition p .

DEFINITION 16. *For p and q in P , we say that $q \leq p$ if*

$$\begin{aligned} \mathcal{S}^p &\subseteq \mathcal{S}^q, \\ \mathcal{S}^p \cap ((\lambda^p + 1) \times \kappa) &= \mathcal{S}^q \cap ((\lambda^p + 1) \times \kappa), \\ \prec^p &= \prec^q \upharpoonright \mathcal{S}^p, \\ \forall x \prec^p y \in \mathcal{S}^p &(\pi_{xy}^p = \pi_{xy}^q), \end{aligned}$$

and λ^p is a mangal of q .

Notice in particular that a condition p may be extended to a condition q simply by extending S^p and f^p — indeed, this is the only way to extend p without changing λ . This kind of extension is analogous to the possibility in Velleman's partial order of extending p by increasing θ_{λ^p} , but we avoid the necessity of (and indeed rule out) adding edges to the tree below level λ^p . We shall sometimes refer to the requirement that λ^p be a mangal of q (which is what is new in this forcing) as the *mangal requirement*; imposing it will ensure that λ^p is a mangal of the generic morass if p is in the generic, as we shall see below.

The only slight difficulty in seeing that \leq as defined above is transitive is with the mangal requirement — whether λ^p is a mangal of r if $p \leq q \leq r$. This is easily resolved, however:

LEMMA 17. *The relation \leq as given in Definition 16 is transitive.*

PROOF. Suppose we have three conditions p , q and r in P_κ such that $p \leq q \leq r$; we wish to show that λ^p is a mangal of r . So suppose there are x and y in r such that $x \prec^r y$ and $l(x) < \lambda^p < l(y)$. If $l(y) \leq \lambda^q$, then $y \in q$ and so since λ^p is a mangal of q , there is a $z \in q$ (and hence $z \in r$ also) such that $l(z) = \lambda^p$ and $x \prec^q z \prec^q y$, whence $x \prec^r z \prec^r y$, as required.

If $l(y) > \lambda^q$, then because λ^q is a mangal of r , there is some $z' \in r$ such that $l(z') = \lambda^q$ and $x \prec^r z' \prec^r y$. But now we are reduced to the previous situation with z' replacing y , and so there is a $z \in r$ with $l(z) = \lambda^p$ and $x \prec^r z \prec^r z'$. Transitivity of \prec^r then gives $x \prec^r z \prec^r y$. \dashv

Thus, letting $\mathbb{P}_\kappa = \langle P_\kappa, \leq \rangle$, we have that \mathbb{P}_κ is a partial order — the *Mangrove Partial Order* or *Mangrove Forcing* at κ . In this section (Section 4) we will generally omit the subscript κ , as κ will be unvarying. Moreover, when we say that morass conditions are compatible or incompatible, or comparable or incomparable, we will mean with respect to this mangrove partial order, rather than Velleman’s morass partial order. We will also abuse notation, writing $p \in \mathbb{P}$ for $p \in P_\kappa$. The goal of the section, realised in Theorem 29, is to show that forcing with \mathbb{P} yields a mangrove in the generic extension.

4.2. A first “bamboo construction”. It is not yet clear that there are any elements of \mathbb{P} other than $\mathbf{1}$. The following lemma constructs one, and the basic technique of the construction, modified from one given in [12], will be reused multiple times as we wish to extend conditions in various ways later on. We call such a construction a “bamboo construction” because of the straight, vertical, unbranching branches that it produces.

LEMMA 18. *There is a morass condition q with $S^q = \omega$.*

PROOF. We start by defining θ_α^q for α up to ω^ω (where the exponentiation is as ordinals), which will be our λ^q . Let

$$\theta_\alpha = \begin{cases} \{n \mid \exists \zeta (\alpha = \omega^n \cdot \zeta)\} & \text{if } \alpha < \omega^\omega \\ \omega & \text{if } \alpha = \omega^\omega \end{cases}$$

Now let

$$\mathcal{S} = (\{\kappa\} \times \omega) \cup \bigcup_{\alpha \leq \omega^\omega} (\{\alpha\} \times \theta_\alpha),$$

$$x \prec^q y \iff l(x) < l(y) \wedge o(x) = o(y),$$

and for $x \prec^q y$ and all $\gamma \leq o(x)$,

$$\pi_{xy}^q(\gamma) = \gamma.$$

We claim that q so defined is a morass condition. The set \mathcal{S} clearly has the right form, \prec^q is a (non-branching!) tree order, and for $x \prec^q y$, π_{xy}^q is a function with the right domain and range. Requirements 1 and 2 for a morass condition are immediate from the construction, as are Monotonicity, Commutativity, M.1, M.2, M.5, M.6 and M.7, the latter two being vacuous in the present context. Finally, axioms M.3 and M.4 below level κ reduce to simple properties of ordinal arithmetic: respectively, that the limit of ordinals divisible by ω^n is divisible by ω^n ; and that if γ is divisible by ω^m for some $m > n$, then the set of ordinals in γ divisible by ω^n is unbounded in γ . \dashv

The reader should note that such a bamboo construction does not reflect the frequent branching that must occur in a mangrove to get κ^+ -many leaves at level κ . However, while one might not expect to find bamboo in a mangrove, it does make a convenient building material.

4.3. μ -equivalence. To acquaint the reader better with morass conditions and mangrove forcing, we now present some lemmas that may help with intuition and which will be useful later on. First, we define a “sameness” notion which appears frequently, and which will be crucial for the homogeneity arguments of Section 6.

DEFINITION 19. *Two morass conditions p and q are μ -equivalent, $p \stackrel{\mu}{\sim} q$, if they are “the same up to level μ ”, that is,*

$$\begin{aligned} \mathcal{S}^p \cap ((\mu + 1) \times \kappa) &= \mathcal{S}^q \cap ((\mu + 1) \times \kappa), \\ \preceq^p \cap ((\mu + 1) \times \kappa)^2 &= \preceq^q \cap ((\mu + 1) \times \kappa)^2, \end{aligned}$$

for $x \preceq^p y$ with $l(y) \leq \mu$,

$$\pi_{xy}^p = \pi_{xy}^q,$$

and μ is a mangal of both p and q .

Note that we include level μ in “up to level μ .” Once again we shall refer to the last requirement in this definition as the mangal requirement.

Clearly $\stackrel{\mu}{\sim}$ is an equivalence relation on the set of those conditions with $\lambda \geq \mu$ and μ as a mangal (although it is not even reflexive for other conditions). While the presence of the mangal requirement means that $\mu \leq \nu$ does not yield $p \stackrel{\nu}{\sim} q \rightarrow p \stackrel{\mu}{\sim} q$, we do have the following.

LEMMA 20. *If $\mu \leq \nu$ and $p \stackrel{\mu}{\sim} q$ and $q \stackrel{\nu}{\sim} r$ for some elements p, q, r of \mathbb{P} , then $p \stackrel{\mu}{\sim} r$.*

PROOF. This is essentially the same as Lemma 17, showing that the \leq relation of \mathbb{P} is transitive. Again, the only difficulty is in showing that μ is a mangal of r . Suppose $x, y \in r$ are such that $x \preceq^r y$ and $l(x) < \mu < l(y)$. If $l(y) \leq \nu$, then $y \in q$ with $x \preceq^q y$. Thus, since μ is a mangal of q , there is a $z \in q$ with $l(z) = \mu$ and $x \preceq^q z \preceq^q y$. So $z \in r$ and $x \preceq^r z \preceq^r y$.

If $l(y) > \nu$, then since ν is a mangal of r , there is a $z \in r$ with $l(z) = \nu$ and $x \preceq^r z \preceq^r y$. By transitivity of \preceq^r , we are reduced to the case above. \dashv

Also, it is immediate from the definitions that $q \leq p \rightarrow q \stackrel{\lambda^p}{\sim} p$. A stronger statement is in fact true.

LEMMA 21. *Let $p, q \in \mathbb{P}$ be (mangrove-) compatible morass conditions, with $\lambda^p \leq \lambda^q$. Then $p \stackrel{\lambda^p}{\sim} q$.*

PROOF. Let $r \in \mathbb{P}$ be such that $r \leq p$ and $r \leq q$. It is immediate from the fact that $p \stackrel{\lambda^p}{\sim} r$ and $q \stackrel{\lambda^q}{\sim} r$ that the necessary equalities for $p \stackrel{\lambda^p}{\sim} q$ hold. To see that λ^p is a mangal of q , observe that since λ^p is a mangal of r , we have that for any x and y in q with $l(x) < \lambda^p < l(y)$,

$$\begin{aligned} x \lhd^q y &\longleftrightarrow x \lhd^r y \\ &\longleftrightarrow \exists z \in r(l(z) = \lambda^p \wedge x \lhd^r z \lhd^r y) \\ &\longleftrightarrow \exists z \in q(l(z) = \lambda^p \wedge x \lhd^q z \lhd^q y), \end{aligned}$$

so λ^p is also a mangal of q . \dashv

Note that, in the situation of the lemma, the only possible obstruction to $q \leq p$ holding is that f^p might not factor through f^q . Indeed, S^p need not be a subset of S^q , in which case f^p certainly cannot factor through f^q .

One reason that the notion of μ -equivalent is important, particularly in the case where $\mu = \lambda^p$ for some condition p , is that most of the “morass-like” requirements for the condition p only pertain to the part of the condition below λ^p . The following lemma makes this observation concrete.

LEMMA 22. *Let p be a morass condition, and let S' be a subset of κ^+ containing 0, closed under ordinal successors and predecessors, and such that $\text{ot}(S') \leq \theta_{\lambda^p}^p$. Then there is a unique morass condition q such that $\lambda^q = \lambda^p$, $q \stackrel{\lambda^p}{\sim} p$, and $S^q = S'$.*

PROOF. The description of q in the statement of the lemma completely determines q : to get from p to q we simply change S^p to S' , correspondingly replace f^p with the order-preserving bijection $f^q : \text{ot}(S') \rightarrow S'$, and modify \lhd and π as appropriate for the new f . To be precise,

$$\mathcal{S}^q = (\mathcal{S}^p \cap ((\lambda^p + 1) \times \kappa)) \cup (\{\kappa\} \times S'),$$

$$\begin{aligned} x \lhd^q y &\longleftrightarrow (l(x) < l(y) \leq \lambda^p \wedge x \lhd^p y) \vee \\ &\quad (l(x) = \lambda^p \wedge l(y) = \kappa \wedge f^q(o(x)) = o(y)) \vee \\ &\quad (l(x) < \lambda^p \wedge l(y) = \kappa \wedge \exists z \in \mathcal{S}^q(l(z) = \lambda^p \wedge x \lhd^q z \lhd^q y)), \end{aligned}$$

and for $x, y \in \mathcal{S}^q$ such that $x \lhd^q y$,

$$\pi_{xy}^q = \begin{cases} \pi_{xy}^p & \text{if } l(x) < l(y) \leq \lambda^p \\ f^q \upharpoonright (o(x) + 1) & \text{if } l(x) = \lambda^p \text{ and } l(y) = \kappa \\ \pi_{zy}^q \circ \pi_{xz}^q & \text{if } l(x) < \lambda^p, l(y) = \kappa, l(z) = \lambda^p \text{ and } x \lhd^q z \lhd^q y. \end{cases}$$

Every aspect of this definition is clearly necessary for q to be a morass condition of the desired form, giving the claimed uniqueness. If we can further show that this definition is sufficient to make q a morass condition we will be done. But now the restricted forms of M.3–M.7 that are required all hold because they do in p (recall from the

discussion after Definition 14 that M.3 and M.5 need only be checked up to level λ). The rest of the requirements for a morass condition are almost all immediate from the definitions; the only points worth mentioning are that f^q is a SLOOP function, and that M.2 in the case that $l(x) < \lambda^p < l(y) = \kappa$ follows from Commutativity and M.2 in the other two cases of the definition of π_{xy}^q . \dashv

In general this q and the original p will not be compatible — if there is a $\tau < \kappa^+$ and $\alpha \neq \beta < \theta_{\lambda^p}^p$ such that $f^p(\alpha) = \tau = f^q(\beta)$, there can be no common extension r , as it would need to satisfy both $\langle \lambda^p, \alpha \rangle \prec^r \langle \kappa, \tau \rangle$ and $\langle \lambda^p, \beta \rangle \prec^r \langle \kappa, \tau \rangle$, violating the fact that \prec^r is a tree relation and Monotonicity holds in r . On the other hand, in Lemma 25 below we shall consider a particular circumstance in which such τ, α and β do not exist, and p and q can be shown to be compatible.

4.4. Cardinal preservation. Our claim is that forcing with \mathbb{P} will yield a mangrove with height κ , with of course $\theta_\kappa = \kappa^+$. For this it is important that the cardinals κ and κ^+ are preserved. In fact, if we assume that we have the GCH in our original model, all cardinals will be preserved, thanks to closure and chain condition properties of \mathbb{P} which we describe here.

Recall that a partial order P is κ -closed if every descending chain of conditions from P of length less than κ has a lower bound in P . Further recall that if P is κ -closed, then forcing with \mathbb{P} adds no new sequences of length less than κ , and so in particular all cardinals $\leq \kappa$ are preserved. In our particular circumstance, knowing that \mathbb{P} is κ -closed will also be useful for “gluing together” successive extensions, making it possible to show that a variety of subsets of \mathbb{P} are dense using just a few basic extension lemmas repeatedly.

One may show that \mathbb{P} is κ -closed by the natural argument, taking the union of the given chain of conditions, and adding a top level as appropriate for the new κ -th level. As pointed out in [12] though, the same argument actually yields a stronger property for \mathbb{P} , one which will be useful when showing later that morasses give rise to mangroves. Perhaps more significantly, it will be exactly what we require in order to obtain master conditions when we force to obtain mangroves while preserving large cardinals.

We make a preliminary definition in order to give Proposition 24 the necessary strength for the argument of Section 5.

DEFINITION 23. *For any $\alpha \leq \kappa^+$, $\mathbb{P}_{\kappa, \alpha}$ denotes the suborder of \mathbb{P} consisting of those conditions $p \in \mathbb{P}$ such that $S^p \subseteq \alpha$.*

One might wonder whether the inclusion $\mathbb{P}_{\kappa, \alpha} \hookrightarrow \mathbb{P}$ is a complete embedding, as defined for example in [10], page 218. It turns out that this depends on α — see Proposition 36 in Section 5.1.

Recall that a set Y in a partial order is *directed* if for every p and q in Y , there is an r in Y such that $r \leq p$ and $r \leq q$. Recall further that a partial order P is κ -directed-closed if every directed subset

Y of P with cardinality less than κ has a lower bound in P . Note in particular that the κ in “ κ -directed-closed” refers to the level of closure rather than the level of directedness. Of course, κ -directed-closure implies κ -closure.

PROPOSITION 24. *For every ordinal $\alpha \leq \kappa^+$, the poset $\mathbb{P}_{\kappa,\alpha}$ is κ -directed-closed. In particular, $\mathbb{P} = \mathbb{P}_{\kappa,\kappa^+}$ is κ -closed, and forcing with \mathbb{P} adds no new sequences of length less than κ of elements of V .*

PROOF. Suppose that Y is a directed set in $\mathbb{P}_{\kappa,\alpha}$ with $|Y| = \gamma$ for some cardinal $\gamma < \kappa$. Let \bar{q} be the direct limit of these structures — that is, let $\mathcal{S}^{\bar{q}} = \bigcup_{p \in Y} \mathcal{S}^p$, $x \prec^{\bar{q}} y$ if and only if for some $p \in Y$ containing both x and y , $x \prec^p y$, and in this case $\pi_{xy}^{\bar{q}} = \pi_{xy}^p$. Note that \bar{q} is well defined: if $p, r \in Y$ both contain x and y , then a common extension $s \in Y$ will also contain x and y , and by the definition of \leq in \mathbb{P} , p and r must both agree with s regarding whether $x \prec y$, and on the function π_{xy} when $x \prec y$ does hold.

Suppose that \bar{q} has a non-empty top level $\lambda^{\bar{q}}$. Then there exists at least one condition p in Y such that $\lambda^p = \lambda^{\bar{q}}$. Recall that the only possibility for extending a condition p without altering λ is to end extend S^p and f^p . Since Y is directed, it follows that conditions p and r in Y with $\lambda^p = \lambda^r = \lambda^{\bar{q}}$ must be comparable: an $s \in Y$ extending both simply has an S^s end-extended from each of S^p and S^r , and so the larger of the latter two sets must simply end-extend the smaller. Thus, the subset $Y_{\lambda^{\bar{q}}}$ of Y consisting of all elements p of Y with $\lambda^p = \lambda^{\bar{q}}$ forms a decreasing chain. Moreover, every element of Y is extended by an element of $Y_{\lambda^{\bar{q}}}$, so \bar{q} is the direct limit of the chain $Y_{\lambda^{\bar{q}}}$. In this case it is straightforward to verify that all of the requirements for a morass condition hold for \bar{q} , simply because they do for each $p \in Y_{\lambda^{\bar{q}}}$. Similarly, it is immediate from the definitions that \bar{q} is an extension of each $p \in Y_{\lambda^{\bar{q}}}$, and hence of each $p \in Y$. Finally, since we have simply taken a union to obtain $\mathcal{S}^{\bar{q}}$, $\mathcal{S}^{\bar{q}} \subseteq \alpha$ and \bar{q} is a lower bound for Y in $\mathbb{P}_{\kappa,\alpha}$.

So suppose now that \bar{q} does not have a top level; then it is not a morass condition and we must modify it to obtain one. Letting $\lambda^q = \sup_{p \in Y}(\lambda^p)$, $S^q = \bigcup_{p \in Y} S^p = \{x \in \mathcal{S}^{\bar{q}} \mid l(x) = \kappa\}$, and f^q be the order preserving bijection from $\text{ot}(S^q)$ to S^q , define q by

$$\mathcal{S}^q = \mathcal{S}^{\bar{q}} \cup (\{\lambda^q\} \times \text{ot}(S^q)),$$

$$\begin{aligned} x \prec^q y \iff & \left(x \prec^{\bar{q}} y \right) \vee \\ & \left(l(x) = \lambda^q \wedge l(y) = \kappa \wedge f^q(o(x)) = o(y) \right) \vee \\ & \left(l(x) < \lambda^q \wedge l(y) = \lambda^q \wedge \right. \\ & \quad \left. \exists z (l(z) = \kappa \wedge x \prec^{\bar{q}} z \wedge f^q(o(y)) = o(z)) \right), \end{aligned}$$

and for $x \prec^q y$,

$$\pi_{xy}^q = \begin{cases} \pi_{xy}^{\bar{q}} & \text{if } x \prec^{\bar{q}} y, \\ f^q \upharpoonright (o(x) + 1) & \text{if } l(x) = \lambda^q \text{ and } l(y) = \kappa, \\ (f^q)^{-1} \circ \pi_{xz}^{\bar{q}} & \text{if } l(y) = \lambda^q, l(z) = \kappa \text{ and } x \prec^{\bar{q}} z. \end{cases}$$

We claim that with this definition q is a morass condition. It is clear that \mathcal{S}^q has the right form, \prec^q is a tree order, and each π_{xy}^q has the right domain and range. Requirement 2 for a morass condition is also immediate from the definition of q . Requirement 1 follows from the fact that it is true for each $p \in Y$ and the easy observation that it also holds at the new top level. Monotonicity, Commutativity and M.1 are similarly straightforward to verify.

For M.2, the cases where both $l(x)$ and $l(y)$ are less than λ^q or where $l(x) < \lambda^q$ and $l(y) = \kappa$ follow from M.2 for the conditions $p \in Y$. If $l(x) = \lambda^q$ and $l(y) = \kappa$, the fact that M.2 is satisfied is immediate from the definition of q . Finally, the case of $l(x) < \lambda^q$ and $l(y) = \lambda^q$ is straightforward from the fact that M.2 is satisfied in some $p \in Y$ containing x and z , where $x \prec^q y \prec^q z$: if $\nu < o(x)$, then

$$\langle l(x), \nu \rangle \prec^p \langle \kappa, \pi_{xz}^p(\nu) \rangle$$

and so

$$\langle l(x), \nu \rangle \prec^q \langle \lambda^q, (f^q)^{-1} \circ \pi_{xz}^p(\nu) \rangle = \langle l(y), \pi_{xy}^q(\nu) \rangle$$

with

$$\pi_{\langle l(x), \nu \rangle \langle l(y), \pi_{xy}^q(\nu) \rangle}^q = (f^q)^{-1} \circ \pi_{xz}^p \upharpoonright (\nu + 1) = \pi_{xy}^q \upharpoonright (\nu + 1)$$

as required.

In the verifications of both M.3 and M.4, the only case we need to consider is when $l(y) = \lambda^q$, since for any $\alpha < \lambda^q$, any condition $p \in Y$ with $\lambda^p \geq \alpha$ fixes the structure of the tree below level α . So suppose that $l(y) = \lambda^q$, and that $y \prec^q z$ with $l(z) = \kappa$. Then there is some $p_z \in Y$ such that $z \in p_z$. For every $r \in Y$ extending p_z ,

$$\begin{aligned} \{l(x) \mid x \prec^q y\} \cap (\lambda^r + 1) &= \{l(x) \mid x \prec^q z\} \cap (\lambda^r + 1) \\ &= \{l(x) \mid x \prec^r z\} \end{aligned}$$

which is closed in $\lambda^r + 1$ by M.3 for r . Moreover, $\lambda^r \in \{l(x) \mid x \prec^r z\}$ for each $r \leq p_z$ in Y since $l(z) = \kappa$. Because $\lambda^q = \sup_{p \in Y}(\lambda^p)$ and Y is directed, we have that the set $\{l(x) \mid x \prec^q y\}$ is closed and unbounded in λ^q , as desired for M.3 and M.4. Also, this implies that for y with $l(y) = \lambda^q$, it is never the case that $x \prec_i^q y$. Hence, the restricted forms of M.6 and M.7 that we require are vacuous at level λ^q , and of course hold below that level because they do in each p_α .

It therefore only remains to check M.5, and again, it is sufficient to consider the case when $l(y) = \lambda^q$. Suppose $\mu < o(y)$; we wish to show that there is an $x \prec^q y$ and a $\nu < o(x)$ such that $\pi_{xy}^q(\nu) = \mu$. Let

$w = \langle \kappa, f^q(\mu) \rangle$, let $z = \langle \kappa, f^q(o(y)) \rangle$ so that $y \prec^q z$, and let $p \in Y$ be such that w and z are both in p . Let $v, x \in p$ with $l(v) = l(x) = \lambda^p$ be such that $x \prec^p z$ and $v \prec^p w$. Then $o(v) < o(x)$, and

$$\pi_{xz}^p(o(v)) = f^p(o(v)) = \pi_{vw}^p(o(v)) = o(w) = f^q(\mu).$$

Now by the definition of \prec^q we have that $x \prec^q y$ and $\pi_{xy}^q(o(v)) = \mu$ as required.

Overall then, we have that q is a morass condition. Moreover, we claim that $q \leq p$ for every $p \in Y$. Indeed, the only part of this claim that is not immediate from the definition of q is the mangal requirement, and even this is trivial to verify when $l(y) \neq \lambda^q$. So suppose $p \in Y$, and $x, y \in q$ with $x \prec^q y$, $l(y) = \lambda^q$ and $l(x) < \lambda^p$. Let $w = \langle \kappa, f^q(o(y)) \rangle$; then $x \prec^q w$. Let $r \in Y$, $r \leq p$ be such that $w \in r$, and hence $x \prec^r w$. By the definition of extension, λ^p is a mangal of r , and so there is a $z \in r$ with $l(z) = \lambda^p$ such that $x \prec^r z \prec^r w$, and hence $x \prec^q z \prec^q y \prec^q w$. Thus, λ^p is a mangal of q for each $p \in Y$, and so $q \leq p$ for each $p \in Y$.

Hence, we have constructed a lower bound q for Y . As before, S^q is simply the union of $\{S^p \mid p \in Y\}$, so $S^q \subseteq \alpha$ and $q \in \mathbb{P}_{\kappa, \alpha}$. Thus, $\mathbb{P}_{\kappa, \alpha}$ is indeed κ -directed-closed. \dashv

Our proof that \mathbb{P} satisfies the κ^+ -cc will require a more complicated construction, building a lower bound for conditions which are the same below level κ but have different sets S at level κ .

LEMMA 25. *Let p and q be morass conditions such that $\lambda^p = \lambda^q$ and $p \lesssim^p q$. Suppose further that there is some $S^0 \subset \kappa^+$ such that both S^p and S^q end-extend S^0 , and $\min(S^q \setminus S^0) \geq \sup(S^p)$. Then p and q are compatible in \mathbb{P} .*

PROOF. We shall build a common extension r of p and q , using a construction similar to that in the proof of Lemma 18.

Let $S^r = S^p \cup S^q$ and let f^r be the order preserving bijection from $\text{ot}(S^r)$ to S^r . Let $\gamma^0 = \text{ot}(S^0)$, $\gamma^p = \text{ot}(S^p)$ and let γ_q be such that $\text{ot}(S^q) = \gamma^0 + \gamma_q$; with this notation, $\text{ot}(S^r) = \gamma^p + \gamma_q$. Note also that by requirements 1 and 2 for morass conditions, γ^0 , γ^p and γ_q are limit ordinals, and $\gamma^p + \gamma_q < \kappa$.

Let $\lambda^r = \lambda^p + \omega^{\gamma^p + \gamma_q}$. For $\alpha \leq \lambda^r$, define θ_α^r by

$$\theta_\alpha^r = \begin{cases} \theta_\alpha^p & \text{if } \alpha \leq \lambda^p \\ \{\eta \mid \exists \zeta (\beta = \omega^\eta \cdot \zeta)\} & \text{if } \alpha = \lambda^p + \beta, 0 < \beta < \omega^{\gamma^p + \gamma_q} \\ \gamma^p + \gamma_q & \text{if } \alpha = \lambda^r. \end{cases}$$

Now, let

$$S^r = (\{\kappa\} \times S^r) \cup \bigcup_{\alpha \leq \lambda^r} (\{\alpha\} \times \theta_\alpha^r),$$

and define \prec^r and the π_{xy}^r (in each case defined only when $x \prec^r y$, even though we do not state that below) according to the following cases. Let id_α denote the identity function on α .

- If $l(x) \geq l(y)$ then $x \not\prec^r y$
- If $l(x) < l(y) \leq \lambda^p$, then $x \prec^r y \leftrightarrow x \prec^p y$, and $\pi_{xy}^r = \pi_{xy}^p$.
- If $\lambda^p < l(x) < l(y) < \kappa$, then $x \prec^r y \leftrightarrow o(x) = o(y)$, and $\pi_{xy}^r = \text{id}_{o(x)+1}$.
- If $\lambda^p = l(x) < l(y) < \kappa$ then
 - if $o(y) < \gamma^p$ then $x \prec^r y \leftrightarrow o(x) = o(y)$ and $\pi_{xy}^r = \text{id}_{o(x)+1}$, and
 - if $o(y) \geq \gamma^p$ then $x \prec^r y \leftrightarrow \exists \zeta (o(x) = \gamma^0 + \zeta \wedge o(y) = \gamma^p + \zeta)$ and

$$\pi_{xy}^r(\alpha) = \begin{cases} \alpha & \text{if } \alpha < \gamma^0 \\ \gamma^p + \eta & \text{if } \alpha = \gamma^0 + \eta. \end{cases}$$

- If $l(x) < \lambda^p < l(y) < \kappa$ then

$$x \prec^r y \leftrightarrow \exists z (l(z) = \lambda^p \wedge x \prec^r z \prec^r y),$$

$$\text{and } \pi_{xy}^r = \pi_{zy}^r \circ \pi_{xz}^r.$$

- If $\lambda^r = l(x) < l(y) = \kappa$ then $x \prec^r y \leftrightarrow f^r(o(x)) = o(y)$ and $\pi_{xy}^r = f^r \upharpoonright (o(x) + 1)$.
- If $l(x) < \lambda^r < l(y) = \kappa$ then

$$x \prec^r y \leftrightarrow \exists z (l(z) = \lambda^r \wedge x \prec^r z \prec^r y),$$

$$\text{and } \pi_{xy}^r = \pi_{zy}^r \circ \pi_{xz}^r.$$

We must verify that r so defined is indeed a morass condition. That \mathcal{S} is of the right form, \prec^r is a tree order and π_{xy}^r is defined where necessary with the right domain and range is clear. Requirements 1 and 2 for a morass condition as well as Monotonicity, Commutativity, M.5, and the restricted form of M.6 required are immediate from the construction; for M.6 note in particular that $\langle \lambda^p, \gamma^0 \rangle$ is \prec^r -less than both $\langle \lambda^p + \omega^{\gamma^p}, \gamma^0 \rangle$ and $\langle \lambda^p + \omega^{\gamma^p}, \gamma^p \rangle$. Once we have observed that Commutativity holds, M.2 also follows, as does M.1 with the earlier observation that γ^p and γ^0 are limit ordinals. Axioms M.3 and M.4 are satisfied as in Lemma 18 along with the fact that they hold in p and hence up to level λ^p in r . Finally, M.7 reduces to a property of ordinal arithmetic: if δ_0 is a limit ordinal, and α is divisible by ω^δ for all $\delta < \delta_0$, then α is divisible by ω^{δ_0} .

Therefore, r is a morass condition. Clearly $r \leq p$ and $r \leq q$; in particular note that the mangal requirement holds by construction. Thus, p and q are compatible. \dashv

PROPOSITION 26. (*GCH*) *The poset \mathbb{P} is κ^+ -cc.*

PROOF. Suppose for the sake of contradiction that A is an antichain in \mathbb{P} of cardinality κ^+ . Since $\lambda^p < \kappa$ for any $p \in \mathbb{P}$, we may assume without loss of generality that $\lambda^p = \lambda$ for all $p \in A$ and some $\lambda < \kappa$. Similarly, there are at most $\kappa^{|\lambda|} = \kappa$ possible functions $\theta : (\lambda + 1) \rightarrow \kappa$ determining the set \mathcal{S} up to level λ , so by further paring down A we may assume that $\mathcal{S}^p \cap ((\lambda + 1) \times \kappa) = \mathcal{S}^q \cap ((\lambda + 1) \times \kappa)$ for all $p, q \in A$. Furthermore, this common initial segment of \mathcal{S} will

have a cardinality $\mu < \kappa$, so there are at most κ many binary relations on it, and we may assume that $\prec^p = \prec^q$ for $p, q \in A$. With \prec fixed, there are at most $(\mu^\mu)^{\mu \times \mu} \leq \kappa$ different possibilities for the set of functions π_{xy} with $l(x) < l(y) \leq \lambda$. Hence, we may assume without loss of generality that the conditions in A are identical up to level λ . Note that this implies that they are completely determined by the set S at level κ : S determines f , which in turn determines the relation $x \prec y$ and maps π_{xy} for $l(y) = \kappa$.

It follows from the Δ -system Lemma (see, for example, [10], Theorem II.1.6) that there is a $B \subset A$ of cardinality κ^+ such that the set $\{S^p \mid p \in B\}$ forms a Δ -system. Moreover, B may be chosen in such a way that there is a $\gamma < \kappa$ and an $S^0 \subset \kappa^+$ such that for all $p, q \in B$, $\text{ot}(S^p) = \text{ot}(S^q) = \gamma$, S^p and S^q end-extend S^0 with $S^p \cap S^q = S^0$, and $\sup(S^p) < \min(S^q \setminus S^0)$ or $\sup(S^q) < \min(S^p \setminus S^0)$. With B so defined, any two of its elements will be compatible by Lemma 25. This contradicts the initial assumption that A was an antichain, and so we may conclude that \mathbb{P} enjoys the κ^+ -cc. \dashv

Note that we have in fact proved the slightly stronger property, that \mathbb{P} is κ^+ -Knaster: for any $A \subset \mathbb{P}$ of cardinality κ^+ , there is a $B \subset A$ of cardinality κ^+ whose elements are pairwise compatible.

4.5. A mangrove from a generic. We now turn to the main task of showing that the direct limit of the conditions in a \mathbb{P} -generic G is a mangrove in $V[G]$. Of course, for this we will need to show that various subsets of \mathbb{P} are dense. The following extension lemmas, in the same vein as Lemma 18 and the argument for Lemma 25, will give us the requisite density.

LEMMA 27. *For any $p \neq \mathbf{1} \in \mathbb{P}$ and any limit ordinal $\sigma < \kappa^+$, there is a $q \leq p$ such that $S^q = S^p \cup ((\sigma + \omega) \setminus \sigma)$.*

PROOF. If $\sigma \in S^p$, then we may of course take $q = p$, so suppose $\sigma \notin S^p$. Let S^q be as in the statement of the lemma. If $\sigma \geq \sup(S^p)$ and $\text{ot}(S^p) + \omega \leq \theta_{\lambda^p}$, we may simply extend p to q by end-extending f^p , making the bijection from $\text{ot}(S^p) + \omega$ to S^q our f^q . Extending as necessary both the tree relation and the maps π_{xy} by composition, it is straightforward to check that this defines a $q \leq p$ with the desired S^q . However, the construction described below works in all cases, so it is not necessary to distinguish this particular case.

If the set $S^p \setminus \sigma$ is non-empty, let τ be its least element, and let $\bar{\tau}$ be the unique ordinal such that $\langle \lambda^p, \bar{\tau} \rangle \prec^p \langle \kappa, \tau \rangle$. Otherwise, let $\bar{\tau} = \text{ot}(S^p)$. Clearly τ (if defined) and $\bar{\tau}$ are limit ordinals. Further, let ξ be such that $\bar{\tau} + \xi = \text{ot}(S^p)$; then ξ is either 0 or a limit ordinal, and $\text{ot}(S^q) = \bar{\tau} + \omega + \xi$.

Note that $\bar{\tau} + \omega + \xi < \kappa$, since $\bar{\tau} + \xi = \text{ot}(S^p) \leq \theta_{\lambda^p} < \kappa$, and so letting $\delta = \omega^{\bar{\tau} + \omega + \xi}$, we also have $\delta < \kappa$. Let $\lambda^q = \lambda^p + \delta$, and let

$$\theta_\alpha^q = \begin{cases} \theta_\alpha^p & \text{if } \alpha \leq \lambda^p \\ \{\nu \mid \exists \zeta(\beta = \omega^\nu \cdot \zeta)\} & \text{if } \alpha = \lambda^p + \beta, 0 < \beta < \delta \\ \bar{\tau} + \omega + \xi & \text{if } \alpha = \lambda^q. \end{cases}$$

Now let

$$\mathcal{S}^q = (\bigcup \{\{\alpha\} \times \theta_\alpha^q \mid \alpha \leq \lambda^q\}) \cup (\{\kappa\} \times S^q);$$

then the usual definitions for θ_α^q and S^q from \mathcal{S}^q match the chosen values, and we also immediately have f^q , defined on the whole width $\theta_{\lambda^q}^q$ of the new top level λ^q .

Let us now define \lhd^q and the functions π_{xy}^q . This will be very similar to the definition for the construction in the argument for Lemma 25. As before, let id_α denote the identity function on α , and although it is not explicitly stated, in each case take π_{xy}^q to be defined only when $x \lhd^q y$.

- If $l(x) \geq l(y)$ then $x \not\lhd^q y$
- If $l(x) < l(y) \leq \lambda^p$, then $x \lhd^q y \leftrightarrow x \lhd^p y$, and $\pi_{xy}^q = \pi_{xy}^p$.
- If $\lambda^p < l(x) < l(y) < \kappa$, then $x \lhd^q y \leftrightarrow o(x) = o(y)$, and $\pi_{xy}^q = \text{id}_{o(x)+1}$.
- If $\lambda^p = l(x) < l(y) < \kappa$ then
 - if $o(y) \leq \bar{\tau}$ then $x \lhd^q y \leftrightarrow o(x) = o(y)$ and $\pi_{xy}^q = \text{id}_{o(x)+1}$, and
 - if $o(y) > \bar{\tau}$ then $x \lhd^q y \leftrightarrow \exists \nu(o(x) = \bar{\tau} + \nu \wedge o(y) = \bar{\tau} + \omega + \nu)$ and

$$\pi_{xy}^q(\alpha) = \begin{cases} \alpha & \text{if } \alpha < \bar{\tau} \\ \bar{\tau} + \omega + \mu & \text{if } \alpha = \bar{\tau} + \mu. \end{cases}$$

- If $l(x) < \lambda^p < l(y) < \kappa$ then

$$x \lhd^q y \leftrightarrow \exists z(l(z) = \lambda^p \wedge x \lhd^q z \lhd^q y),$$

$$\text{and } \pi_{xy}^q = \pi_{zy}^q \circ \pi_{xz}^q.$$

- If $\lambda^q = l(x) < l(y) = \kappa$ then $x \lhd^q y \leftrightarrow f^q(o(x)) = o(y)$ and $\pi_{xy}^q = f^q \upharpoonright (o(x) + 1)$.
- If $l(x) < \lambda^q < l(y) = \kappa$ then

$$x \lhd^q y \leftrightarrow \exists z(l(z) = \lambda^q \wedge x \lhd^q z \lhd^q y),$$

$$\text{and } \pi_{xy}^q = \pi_{zy}^q \circ \pi_{xz}^q.$$

That completes the definition of q . It is now a simple matter to check that the axioms for a morass condition hold for q ; indeed, it is an identical verification process to that for the r of Lemma 25.

So q is a condition with $S^q = S^p \cup ((\sigma + \omega) \setminus \sigma)$, and by construction q extends p . \dashv

LEMMA 28. *For any $p \neq \mathbf{1} \in \mathbb{P}$ there is a $q \leq p$ such that $S^q = S^p$, $\theta_{\lambda^q}^q = \text{ot}(S^q)$, and $\lambda^q = \lambda^p + \omega^{\text{ot}(S^q)}$.*

PROOF. The proof is another variant of the same argument, this one being particularly straightforward. Let S^q and λ^q be as in the statement, set

$$\theta_\alpha^q = \begin{cases} \theta_\alpha^p & \text{if } \alpha \leq \lambda^p \\ \{\nu \mid \exists \zeta (\beta = \omega^\nu \cdot \zeta)\} & \text{if } \alpha = \lambda^p + \beta, 0 < \beta < \omega^{\text{ot}(S^q)} \\ \text{ot}(S^q) & \text{if } \alpha = \lambda^q, \end{cases}$$

and let

$$\mathcal{S}^q = \left(\bigcup \{ \{\alpha\} \times \theta_\alpha^q \mid \alpha \leq \lambda^q \} \right) \cup (\{\kappa\} \times S^q).$$

As always, take f^q to be the order preserving bijection from $\text{ot}(S^q)$ to S^q . Define $x \lhd^q y$, and for x and y such that $x \lhd^q y$, π_{xy}^q , by the following cases. Again, id_α denotes the identity function on α .

- If $l(x) \geq l(y)$ then $x \not\lhd^q y$
- If $l(x) < l(y) \leq \lambda^p$, then $x \lhd^q y \leftrightarrow x \lhd^p y$, and $\pi_{xy}^q = \pi_{xy}^p$.
- If $\lambda^p \leq l(x) < l(y) < \kappa$, then $x \lhd^q y \leftrightarrow o(x) = o(y)$, and $\pi_{xy}^q = \text{id}_{o(x)+1}$.
- If $l(x) < \lambda^p < l(y) < \kappa$ then

$$x \lhd^q y \leftrightarrow \exists z (l(z) = \lambda^p \wedge x \lhd^q z \lhd^q y),$$

$$\text{and } \pi_{xy}^q = \pi_{zy}^q \circ \pi_{xz}^q.$$

- If $\lambda^q = l(x) < l(y) = \kappa$ then $x \lhd^q y \leftrightarrow f^q(o(x)) = o(y)$ and $\pi_{xy}^q = f^q \mid (o(x) + 1)$.
- If $l(x) < \lambda^q < l(y) = \kappa$ then

$$x \lhd^q y \leftrightarrow \exists z (l(z) = \lambda^q \wedge x \lhd^q z \lhd^q y),$$

$$\text{and } \pi_{xy}^q = \pi_{zy}^q \circ \pi_{xz}^q.$$

It is routine to check that with this definition q is a morass condition and $q \leq p$. \dashv

Now let G be \mathbb{P} -generic over V . Let $M^G = \langle \mathcal{S}^G, \lhd^G, \langle \pi_{xy}^G \rangle_{x \lhd^G y} \rangle$ be the direct limit of the system G in $V[G]$. That is, \mathcal{S}^G is the union over $p \in G$ of \mathcal{S}^p , for x and y in \mathcal{S}^G we set $x \lhd^G y$ if and only if there is some condition $p \in G$ containing x and y such that $x \lhd^p y$, and in this case $\pi_{xy}^G = \pi_{xy}^p$.

THEOREM 29. *With M^G defined as above, M^G is a mangrove in $V[G]$.*

PROOF. Clearly \mathcal{S}^G is a subset of $(\kappa \times \kappa) \cup (\{\kappa\} \times \kappa^+)$ and the maps π_{xy}^G are defined when they ought to be with the right domain and range. The relation \lhd^G is tree relation since \lhd^p is a tree relation for each condition p , and below level κ extensions of a condition only end-extend the relation. From Lemma 27, we have that $D_\tau = \{p \in \mathbb{P} \mid \tau \in S^p\}$ is dense in \mathbb{P} for each $\tau < \kappa^+$, and so it follows that $G \cap D_\tau \neq \emptyset$ and hence $S^G = \kappa^+$. By Lemma 28 along with the κ -closure property of \mathbb{P} , it also follows that for any $\alpha < \kappa$, it is dense for the condition p to have $\lambda^p > \alpha$. In particular, θ_α^G is an

ordinal greater than 0 for every $\alpha \leq \kappa$, and the full strength of the Left-alignment axiom is satisfied.

Monotonicity, Commutativity, M.1 and M.2 hold in M^G because they do in each condition. Similarly, to show that M.3 and M.4 hold in M^G , it suffices to show that they do for $y \in M^G$ with level κ . So suppose $y \in M^G$ and $l(y) = \kappa$. Given $\alpha < \kappa$, we may take a condition p such that $y \in p$ and $\lambda^p > \alpha$. Then there is some $z \in p$ with $l(z) = \lambda^p$ such that $z \prec^p y$. Hence, the set $\{l(z) \mid z \prec^G y\}$ is unbounded in κ , as required for M.4. Also, since the corresponding set in each condition is closed in κ , and the top levels λ^p of the conditions are unbounded in κ , we also have M.3: that the set is closed in κ .

Axiom M.5 is also immediate everywhere in M^G except level κ , from the fact that it holds in each condition. So suppose $y \in M^G$ has $l(y) = \kappa$, and let $\alpha < o(y)$. Let $w = \langle \kappa, \alpha \rangle$, and let $p \in G$ be such that $w, y \in p$. Let $v, x \in p$ be such that $l(v) = l(x) = \lambda^p$, $v \prec^p w$ and $x \prec^p y$. Then by the remarks following definition 14, $\pi_{xy}^p(o(v)) = \alpha$. Thus, M.5 holds in M^G at level κ as well.

Axioms M.6 and M.7 require more work to verify, but fortunately the situation is the same as in Velleman's paper [12]. We follow the arguments there, but recast them as transfinite inductions on natural well orders of the elements of M^G , rather than proofs by contradiction for the least counterexample.

To verify M.6, suppose that $x \prec^G y \in M^G$ with $o(x)$ a limit ordinal, and that M.6 holds for all such $x' \prec^G y'$ with y' lexicographically less than y ; that is, $l(y') < l(y)$ or $l(y') = l(y) \wedge o(y') < o(y)$. Let $\zeta = \sup(\pi_{xy}^G "o(x))$ and let $z = \langle l(y), \zeta \rangle$. Clearly if $z = y$ we are done, so suppose $\zeta < o(y)$.

If y is a limit point of \prec^G , then M.5 holds of y and there is some $w \in M^G$ and $\mu < o(w)$ such that $x \prec^G w \prec^G y$ and $\pi_{wy}(\mu) = \zeta$. Denote $\langle l(w), \mu \rangle$ by u ; then by axiom M.2, $u \prec^G z$, and $\pi_{uz}^G = \pi_{wy}^G \upharpoonright (\mu + 1)$. By the composition axiom and M.1, we see that $\pi_{xw}^G "o(x) \subset \mu$, and moreover since $\pi_{xy}^G "o(x)$ is unbounded in $\zeta = \pi_{wy}^G(\mu)$, $\pi_{xw}^G "o(x)$ must be unbounded in μ . But now by the induction hypothesis, $x \prec^G u$ with $\pi_{xu}^G \upharpoonright o(x) = \pi_{xw}^G \upharpoonright o(x)$. Hence composing, we have that $x \prec^G z$ and

$$\pi_{xz}^G \upharpoonright o(x) = \pi_{uz}^G \circ \pi_{xu}^G \upharpoonright o(x) = \pi_{wy}^G \upharpoonright (\mu + 1) \circ \pi_{xw}^G \upharpoonright o(x) = \pi_{xy}^G \upharpoonright o(x)$$

as required.

On the other hand, if y is not a limit point of \prec^G , we may take a $w \in M^G$ such that $w \prec_i^G y$. Note that by M4, $l(y) \neq \kappa$ in this case. If $w = x$ then M.6 for x and y follows from the fact that it holds in any condition in G containing both x and y . So suppose that $x \neq w$ and hence $x \prec^G w \prec_i^G y$. Let $\mu = \sup(\pi_{xw}^G "o(x))$ and $u = \langle l(w), \mu \rangle$, so that by the induction hypothesis, $x \prec^G u$ with $\pi_{xu}^G \upharpoonright o(x) = \pi_{xw}^G \upharpoonright o(x)$. Note that it is possible that $u = w$. Let $\nu = \pi_{wy}(\mu)$ and $v = \langle l(y), \nu \rangle$; then if $\mu < o(w)$, $u \prec^G v$ with

$\pi_{uv}^G = \pi_{wy}^G \upharpoonright (\mu + 1)$ by M.2, and otherwise $u = w$, $v = y$ and the same statement trivially holds true. But now since $\pi_{xw}^G \text{“} o(x) \text{”}$ is by definition unbounded in μ ,

$$\begin{aligned} \sup(\pi_{uv}^G \text{“} \mu \text{”}) &= \sup(\pi_{uv}^G \circ \pi_{xw}^G \text{“} o(x) \text{”}) \\ &= \sup(\pi_{wy}^G \upharpoonright (\mu + 1) \circ \pi_{xw}^G \text{“} o(x) \text{”}) \\ &= \sup(\pi_{xy}^G \text{“} o(x) \text{”}) \\ &= \zeta \end{aligned}$$

If $\mu < o(w)$ and $\nu < o(y)$ we have by the induction hypothesis that $u \prec^G z$ and $\pi_{uz}^G \upharpoonright \mu = \pi_{uv}^G \upharpoonright \mu$. Otherwise, $u = w$ and $v = y$, so that $u \prec_i^G v$ and we also obtain $u \prec^G z$ and $\pi_{uz}^G \upharpoonright \mu = \pi_{uv}^G \upharpoonright \mu$, this time from the fact that it holds in any condition containing both u and v . So then $x \prec^G z$, and

$$\pi_{xz}^G \upharpoonright o(x) = \pi_{uz}^G \circ \pi_{xu}^G \upharpoonright o(x) = \pi_{uv}^G \circ \pi_{xw}^G \upharpoonright o(x) = \pi_{wy}^G \circ \pi_{xw}^G \upharpoonright o(x) = \pi_{xy}^G \upharpoonright o(x).$$

Thus, we have by transfinite induction on the lexicographic order for y that M.6 holds of M^G .

To verify M.7 for M^G we will induct simply on $l(y)$. Let x, y and α be as in the antecedent of M.7, and suppose first that there is some w such that $x \prec^G w \prec^G y$ and $l(w) > \alpha$. This is in particular the case when y is a limit node of the tree, thanks to Monotonicity and M.3. For each $\nu < o(x)$, we have from M.2 that

$$\langle l(x), \nu \rangle \prec^G \langle l(w), \pi_{xw}^G(\nu) \rangle \prec^G \langle l(y), \pi_{wy}^G \circ \pi_{xw}^G(\nu) \rangle = \langle l(y), \pi_{xy}^G(\nu) \rangle.$$

Thus, since $\pi_{xy}^G \text{“} o(x) \text{”}$ is unbounded in $o(y)$, it must be the case that $\pi_{xw}^G \text{“} o(x) \text{”}$ is unbounded in $o(w)$. But now x, w and α fit the hypotheses for M.7, and $l(w) < l(y)$, so by the induction hypothesis, there is a γ such that $\langle \alpha, \gamma \rangle \prec^G w \prec^G y$.

So now suppose we are in the other situation, where there is some w with $l(w) \leq \alpha$ such that $w \prec_i^G y$. If $l(w) = \alpha$ we are clearly done. Also, if w were x , then M.7 would hold for x and y since it does for any condition containing them, and that would contradict the assumption that α is as in the hypothesis for that axiom. So suppose that $x \prec^G w \prec_i^G y$ with $l(w) < \alpha$; we will derive a similar contradiction in this case. As in the above case, we have for each $\nu < o(x)$ that

$$\langle l(x), \nu \rangle \prec^G \langle l(w), \pi_{xw}^G(\nu) \rangle \prec^G \langle l(y), \pi_{xy}^G(\nu) \rangle,$$

and $\pi_{xw}^G \text{“} o(x) \text{”}$ is unbounded in $o(w)$. Let us denote by γ_ν that ordinal such that $\langle \alpha, \gamma_\nu \rangle \prec^G \langle l(y), \pi_{xy}^G(\nu) \rangle$, as we are assuming exists. For simplicity, let us also define notation for $\langle l(x), \nu \rangle$, $\langle l(w), \pi_{xw}^G(\nu) \rangle$, and $\langle l(y), \pi_{xy}^G(\nu) \rangle$ for any given $\nu < o(x)$: call them x_ν , w_ν and y_ν respectively. Then our string of tree relations can be rewritten and extended thus:

$$x_\nu \prec^G w_\nu \prec^G \langle \alpha, \gamma_\nu \rangle \prec^G y_\nu.$$

Now, given $\mu < o(w)$, take some $\nu < o(x)$ such that $\pi_{xw}^G(\nu) > \mu$. By M.2, we have that

$$\langle l(w), \mu \rangle \prec^G \langle \alpha, \pi_{w_\nu \langle \alpha, \gamma_\nu \rangle}^G(\mu) \rangle \prec^G \langle l(y), \pi_{w_\nu y_\nu}^G(\mu) \rangle,$$

and the corresponding functions π are the restriction of $\pi_{w_\nu \langle \alpha, \gamma_\nu \rangle}^G$ to $\mu + 1$ and the restriction of $\pi_{\langle \alpha, \gamma_\nu \rangle y_\nu}^G$ to $\pi_{w_\nu \langle \alpha, \gamma_\nu \rangle}^G(\mu) + 1$. We now have that the hypotheses for M.7 hold of w , y and α , and $w \prec_i^G y$, so the conclusion also obtains, since it does in any condition containing w and y . Of course, an ordinal γ given by M.7 for w , y , and α would also act as a witness to the truth of M.7 for x , y , and α . But in fact, we have a contradiction (as before) to the assumption that $w \prec_i^G y$, and so the supposed situation of having w satisfying $x \prec^G w \prec_i^G y$ and $l(w) < \alpha$ cannot occur. In any case, we may conclude that M.7 holds in M^G .

That completes the verification that M^G is a morass. To confirm that it is a mangrove, note that for any $p \in G$, λ^p is a mangal of M^G : this is immediate from the mangal requirement for extension. Thus, since the λ^p for $p \in G$ are unbounded in κ , M^G is a mangrove, as claimed. \dashv

§5. A mangrove from a morass. In this section we deviate from our main programme in order to answer a natural question that arose earlier — whether the existence of a morass implies the existence of a mangrove. A positive solution is provided by the theorems of [12], in which Velleman proves the equivalence of the existence of a morass with a certain forcing axiom. We shall show that this forcing axiom is sufficient to generate a mangrove from the mangrove forcing \mathbb{P} defined in the previous section. Indeed, our argument will closely follow Velleman’s argument that the forcing axiom gives a morass from his partial order.

We start with some preliminary definitions from [12]. Let P be a partial order, and let $\mathcal{D} = \{D_\alpha \mid \alpha < \kappa^+\}$ be an indexed family of open dense sets in P . For each $p \in P$, define the *realm* of p , $\text{rlm}(p)$, to be the set $\{\alpha < \kappa^+ \mid p \in D_\alpha\}$. For each $\alpha < \kappa^+$ let $P_\alpha = \{p \in P \mid \text{rlm}(p) \subseteq \alpha\}$, and let $P^* = \bigcup_{\alpha < \kappa} P_\alpha$, the set of conditions with realms bounded below κ .

DEFINITION 30 (Definition 1.1.2 of [12]). *A family $\mathcal{D} = \{D_\alpha \mid \alpha < \kappa^+\}$ of subsets of P is said to be almost κ -indiscernible if*

- I.1:** $P^* \neq \emptyset$, and for all $\alpha < \kappa$, $D_\alpha \cap P^*$ is open dense in P^* , and
- I.2:** for all $\alpha < \kappa$, P_α is κ -directed-closed,

and there is a function σ assigning to each SLOOP function $f : \alpha \rightarrow \gamma$ with $\alpha < \kappa$ and $\gamma < \kappa^+$ a function $\sigma_f : P_\alpha \rightarrow P_\gamma$ such that the following properties also hold for all such f .

- I.3:** σ_f is order preserving. That is,

$$\forall p, q \in P_\alpha (p \leq q \rightarrow \sigma_f(p) \leq \sigma_f(q)).$$

- I.4:** For all $p \in P_\alpha$, $\text{rlm}(\sigma_f(p)) = f^{\text{“}}\text{rlm}(p)$.
I.5: If $\gamma < \kappa$, and for some $\beta < \alpha$, $f \restriction \beta = \text{id}_\beta$ and $f(\beta) \geq \alpha$, then for all $p \in P_\alpha$, p and $\sigma_f(p)$ are compatible in P^* .
I.6: If $f_1 : \alpha_1 \rightarrow \alpha_2$ and $f_2 : \alpha_2 \rightarrow \gamma$ are strictly order preserving with $\alpha_1, \alpha_2 < \kappa$ and $\gamma < \kappa^+$, then

$$\sigma_{f_2 \circ f_1} = \sigma_{f_2} \circ \sigma_{f_1}.$$

One of the main results of [12] is the following.

THEOREM 31 (Theorems 1.1.3 and 2.1.6 of [12]). *The following are equivalent:*

1. *There exists a $(\kappa, 1)$ -morass.*
2. *For any partial order P and \mathcal{D} an almost κ -indiscernible family of subsets of P , there is a filter G in P such that for every $D_\alpha \in \mathcal{D}$, $G \cap D_\alpha \neq \emptyset$.*

Velleman further strengthens (2) above by making G hit κ -many additional sets E_ζ ; the reader trying to reconcile the statements here and in [12] may take each E_ζ to be P . In this way statement (2) above can be seen to lie in strength between the formulations in Theorems 1.1.3 and 2.1.6 of [12], which are shown there to actually be equivalent.

To show that the existence of a morass implies the existence of a mangrove, we will show that the subsets of the mangrove forcing

$$D_\alpha = \{p \in \mathbb{P}_\kappa \mid \alpha \in S^p\}$$

for $\alpha < \kappa^+$ form an almost κ -indiscernible family. The filter we then obtain from Theorem 31 will have a mangrove as its direct limit, using essentially the same proof as for Theorem 29, where the filter in question was \mathbb{P} -generic over V .

Observe that with these sets D_α , $\text{rlm}(p) = S^p$ for all $p \in \mathbb{P}_\kappa$, and P_α as defined before Definition 30 is simply the set of all morass conditions p with $S^p \subseteq \alpha$. That is, $P_\alpha = \mathbb{P}_{\kappa, \alpha}$ as in Definition 23; we shall stick with the $\mathbb{P}_{\kappa, \alpha}$ notation here. Also, recall that for any morass condition p , $|S^p| \leq \theta_{\lambda^p} < \kappa$, so if $p \in \mathbb{P}_{\kappa, \kappa}$, then there is some $\alpha < \kappa$ such that $p \in \mathbb{P}_{\kappa, \alpha}$. Thus, $P^* = \mathbb{P}_{\kappa, \kappa}$; again, we will stick with the notation from Definition 23.

PROPOSITION 32. *The sequence $\mathcal{D} = \langle D_\alpha \mid \alpha < \kappa^+ \rangle$ as defined above is an almost κ -indiscernible family of subsets of the mangrove forcing \mathbb{P}_κ .*

PROOF. Lemma 18 shows that $\mathbb{P}_{\kappa, \kappa}$ is non-empty, and Lemma 27 shows that for each $\alpha < \kappa$, $D_\alpha \cap \mathbb{P}_{\kappa, \kappa}$ is dense, with openness being clear from the definition of \leq in \mathbb{P} . Therefore, I.1 holds. Also, I.2 is just Proposition 24 for $\alpha < \kappa$.

We must now define $\sigma_f : \mathbb{P}_{\kappa, \alpha} \rightarrow \mathbb{P}_{\kappa, \gamma}$ for f a SLOOP function with domain $\alpha < \kappa$ and codomain $\gamma < \kappa^+$. If $p \in \mathbb{P}_{\kappa, \alpha}$, let $\sigma_f(p)$ be that $q \in \mathbb{P}_{\kappa, \gamma}$ such that $\lambda^q = \lambda^p$, $q \stackrel{\lambda^p}{\sim} p$, and $S^q = f^{\text{“}}S^p$, as given by

Lemma 22. Of course, this definition makes I.4 and I.6 immediate. Further, the antecedent of I.5 makes Lemma 25 applicable, and the common extension constructed in that lemma clearly lies in $\mathbb{P}_{\kappa, \kappa}$, so I.5 also holds.

To check I.3, let g be a SLOOP function, and suppose $p \leq q$. Clearly λ^q remains a mangal of $\sigma_g(p)$, and we have $\sigma_g(q) \stackrel{\lambda^q}{\sim} \sigma_g(p)$. If we can now show that

$$\begin{aligned} \Vdash^{\sigma_g(q)} \upharpoonright ((\{\lambda^q\} \times \theta_{\lambda^q}^{\sigma_g(q)}) \times (\{\kappa\} \times S^{\sigma_g(q)})) \\ = \Vdash^{\sigma_g(p)} \upharpoonright ((\{\lambda^q\} \times \theta_{\lambda^q}^{\sigma_g(q)}) \times (\{\kappa\} \times S^{\sigma_g(q)})), \end{aligned}$$

and that for $x \Vdash^{\sigma_g(q)} y \in \mathcal{S}^{\sigma_g(q)}$ with $l(x) = \lambda^q$ and $l(y) = \kappa$, $\pi_{xy}^{\sigma_g(q)} = \pi_{xy}^{\sigma_g(p)}$, then we will be done, by Commutativity and the fact that $\lambda^q = \lambda^{\sigma_g(q)}$ is a mangal of $\sigma_g(q)$. Further, since $f^{\sigma_g(q)}$ is surjective onto $S^{\sigma_g(q)}$, it is sufficient to check that $\Vdash^{\sigma_g(q)} \subseteq \Vdash^{\sigma_g(p)}$ and $\pi_{xy}^{\sigma_g(q)} = \pi_{xy}^{\sigma_g(p)}$ on the relevant domain.

So suppose that $x \Vdash^{\sigma_g(q)} y$ for some $x, y \in \sigma_g(q)$ with $l(x) = \lambda^q$ and $l(y) = \kappa$. Note that by the definition of $\sigma_g(q)$, $f^{\sigma_g(q)} = g \circ f^q$, and likewise with q replaced by p . Now

$$x \Vdash^q \langle \kappa, f^q(o(x)) \rangle,$$

so

$$x \Vdash^p \langle \kappa, f^q(o(x)) \rangle.$$

Let z be that $z \in p$ such that $l(z) = \lambda^p$ and

$$x \Vdash^p z \Vdash^p \langle \kappa, f^q(o(x)) \rangle.$$

Then

$$x \Vdash^{\sigma_g(p)} z \Vdash^{\sigma_g(p)} \langle \kappa, g \circ f^q(o(x)) \rangle = \langle \kappa, f^{\sigma_g(q)}(o(x)) \rangle = y$$

as required.

Similarly,

$$\begin{aligned} \pi_{xy}^{\sigma_g(q)} &= f^{\sigma_g(q)} \upharpoonright (o(x) + 1) \\ &= g \circ f^q \upharpoonright (o(x) + 1) \\ &= g \circ \pi_{xy}^q \\ &= g \circ \pi_{xy}^p \\ &= g \circ f^p \circ \pi_{xz}^p \quad \text{for } z \text{ as above} \\ &= f^{\sigma_g(p)} \circ \pi_{xz}^p \\ &= f^{\sigma_g(p)} \circ \pi_{xz}^{\sigma_g(p)} \\ &= \pi_{xy}^{\sigma_g(p)} \end{aligned}$$

and we are done. \dashv

From Proposition 32 and Theorem 31, we obtain that if a morass exists at κ , then there is a filter G in \mathbb{P}_κ intersecting every D_α .

In the proof of Theorem 29, the only dense sets G was required to meet were the sets D_α , along with the open dense set of conditions

p with $\lambda^p > \gamma$, for each $\gamma < \kappa$. It turns out that using the following argument from [12] we can avoid directly requiring G to meet these latter sets.

LEMMA 33. *If G is a filter over \mathbb{P} such that $G \cap D_\alpha \neq \emptyset$ for each $\alpha < \kappa^+$, then $\{\lambda^p \mid p \in G\}$ is cofinal in κ .*

PROOF. It will be convenient to work with the direct limit M^G of the filter G , with components \mathcal{S}^G , \prec^G and π^G . Observe first that the verification of M.5 at level κ in M^G in the proof of Theorem 29 only required that G intersect every D_α , and thus goes through in our situation. We also have that in M^G , \prec is a tree order and Monotonicity holds, since these properties are true in every condition, and below level κ the relation \prec^p is only end-extended in moving to an extension of p . In particular, we have that for every $y \in \mathcal{S}^G$, $\{x \mid x \prec y\}$ contains at most one element from each level. Consider the node $y = \langle \kappa, \kappa \rangle$ in M^G . For every $x \prec^G y$, $|\pi_{xy}^G \text{``} o(x) \text{"}| = |o(x)| < \kappa$. Thus, since κ is regular, there must be κ distinct $x \prec^G y$, and $\{l(x) \mid x \prec^G y\}$ is cofinal in κ . But now for each $x \in \mathcal{S}^G$, there is a $p \in G$ such that $x \in p$, and hence $l(x) \leq \lambda^p$. It follows that $\{\lambda^p \mid p \in G\}$ is cofinal in κ . \dashv

THEOREM 34. *There is a morass at κ if and only if there is a mangrove at κ .*

PROOF. Suppose there is a morass at κ . As mentioned above, by proposition 32 and Theorem 31, there is a filter G in \mathbb{P}_κ intersecting every D_α . By Lemma 33, $\{\lambda^p \mid p \in G\}$ is cofinal in κ . Knowing this, the proof of Theorem 29 goes through for G , giving that the direct limit M^G is a mangrove. Of course, the other direction is trivial. \dashv

5.1. Aside: Complete embeddings. Although it is not necessary for the preceding arguments, the curious reader might wonder whether the inclusion maps $\mathbb{P}_{\kappa,\alpha} \hookrightarrow \mathbb{P}_\kappa$ are complete embeddings. This seemingly simple question actually takes quite some work to answer, so we here devote a subsection to its resolution.

Recall the following definition from [10].

DEFINITION 35. *Let P and Q be partial orders. A function $i : P \rightarrow Q$ is a complete embedding if*

1. $\forall p, p' \in P (p' \leq p \rightarrow i(p') \leq i(p))$
2. $\forall p, p' \in P (p \perp p' \leftrightarrow i(p) \perp i(p'))$
3. $\forall q \in Q \exists p \in P \forall p' \in P (p' \leq p \rightarrow i(p') \parallel q)$

A condition p as in 3 is said to be a reduction of q to P .

PROPOSITION 36. *For any limit ordinal $\alpha \leq \kappa^+$, the inclusion $\mathbb{P}_{\kappa,\alpha} \hookrightarrow \mathbb{P}_\kappa$ is a complete embedding if and only if $\text{cf}(\alpha) = \kappa$.*

Note that because of the requirement on morass conditions that S be closed under ordinal successors, $\mathbb{P}_{\kappa,\alpha+n} = \mathbb{P}_{\kappa,\alpha}$ for any limit ordinal α and natural number n . The proposition therefore effectively covers all cases.

PROOF. Suppose first that $\alpha < \kappa$. Let $q \in \mathbb{P}_\kappa$ be a morass condition such that $\text{ot}(S^q) > \alpha$ (such a q may be constructed using lemmas 18 and 27 and Proposition 24). Let $p \in \mathbb{P}_{\kappa,\alpha}$ be arbitrary; we shall show that p is not a reduction of q to $\mathbb{P}_{\kappa,\alpha}$.

Since $p \in \mathbb{P}_{\kappa,\alpha}$, $\text{ot}(S^p) \leq \alpha$. Note that in the constructions for Lemma 28 and Proposition 24, the extension q obtained satisfies $\theta_{\lambda^q} = \text{ot}(S^q)$. Thus, applying these results, we have that we can extend p to a $p' \in \mathbb{P}_{\kappa,\alpha}$ with $\lambda^{p'} > \lambda^q$ and $\theta_{\lambda^{p'}}^{p'} \leq \alpha$. We claim that p' is incompatible with q : if there were an $r \in \mathbb{P}_\kappa$ extending both p' and q , then $\lambda^{p'}$ would be a mangal of r , with $\theta_{\lambda^{p'}}^r = \theta_{\lambda^{p'}}^{p'} \leq \alpha$. But the edge $\langle \lambda^q, \alpha \rangle \prec^q \langle \kappa, f^q(\alpha) \rangle$ of q , and hence of r , cannot factor through such a level $\lambda^{p'}$ — as a consequence of M.1, $o(x) \leq o(y)$ for any $x \prec^r y$. Thus, no such common extension r can exist, and p is not a reduction of q to $\mathbb{P}_{\kappa,\alpha}$. Therefore the inclusion $\mathbb{P}_{\kappa,\alpha} \hookrightarrow \mathbb{P}_\kappa$ is not a complete embedding.

Suppose next that $\text{cf}(\alpha) = \kappa$, and let q be any element of $\mathbb{P} \setminus \mathbb{P}_{\kappa,\alpha}$. Let $S_\alpha^q = S^q \cap \alpha$; because $\text{ot}(S_\alpha^q) \leq \text{ot}(S^q) < \kappa$, S_α^q is bounded in α , and we will be able to overcome the problems seen in the $\alpha < \kappa$ case. Let $\tau = \sup(S_\alpha^q)$ and $\mu = \text{ot}(S^q \setminus S_\alpha^q)$. Using Lemma 22, let r be the condition such that $\lambda^r = \lambda^q$, $r \stackrel{\lambda^q}{\sim} q$, and $S^r = S_\alpha^q \cup ((\tau + \mu) \setminus \tau)$. We claim that r is a reduction of q to $\mathbb{P}_{\kappa,\alpha}$.

Suppose $p \leq r$ in $\mathbb{P}_{\kappa,\alpha}$, and note in particular that this implies that $S^p \supseteq S^r$. Applying Lemma 22 again, we can obtain a $p' \in \mathbb{P}_\kappa$ with $\lambda^{p'} = \lambda^p$, $p' \stackrel{\lambda^p}{\sim} p$, and $S^{p'} = (S^p \cap \tau) \cup (S^q \setminus S_\alpha^q)$. Applying the construction of Lemma 25, we obtain a condition s extending both p and p' , with $S^s = S^p \cup (S^q \setminus S_\alpha^q)$. We claim that $s \leq q$. Certainly S^s , and in particular S^s , is large enough. We also have that

$$q \stackrel{\lambda^q}{\sim} r \stackrel{\lambda^r}{\sim} p \stackrel{\lambda^p}{\sim} s$$

so $q \stackrel{\lambda^q}{\sim} s$. Since λ^q is a mangal of q , it suffices now to check that

$$\prec^q \upharpoonright ((\{\lambda^q\} \times \theta_{\lambda^q}^q) \times (\{\kappa\} \times S^q)) = \prec^s \upharpoonright ((\{\lambda^q\} \times \theta_{\lambda^q}^q) \times (\{\kappa\} \times S^q)),$$

and that for $x \prec^q y \in S^q$ with $l(x) = \lambda^q$ and $l(y) = \kappa$, $\pi_{xy}^q = \pi_{xy}^s$.

Now for x and y in q with $l(x) = \lambda^q$ and $l(y) = \kappa$, $x \prec^q y$ if and only if $f^q(o(x)) = o(y)$, and $\pi_{xy}^q = f^q \upharpoonright (o(x) + 1)$. It therefore suffices to show that for such x and y , $x \prec^s y$ whenever $f^q(o(x)) = o(y)$, with $\pi_{xy}^s = f^q \upharpoonright (o(x) + 1)$ — the converse holds because f^q , the order preserving bijection from $\text{ot}(S^q)$ to S^q , is surjective onto S^q . So suppose $x \prec^q y$, that is, $f^q(o(x)) = o(y)$. There are two cases to consider.

- If $o(x) < \text{ot}(S_\alpha^q)$, then x and y are actually in r , and we have $x \prec^r y$ with $\pi_{xy}^r = \pi_{xy}^q$. Since $r \geq p \geq s$, we have $x \prec^s y$ with $\pi_{xy}^s = \pi_{xy}^q$.

- If $\text{ot}(S_\alpha^q) \leq o(x) < \text{ot}(S^q)$, let β be such that $o(x) = \text{ot}(S_\alpha^q) + \beta$. Then in r ,

$$x \Vdash^r \langle \kappa, \tau + \beta \rangle, \text{ with } \pi_{x \langle \kappa, \tau + \beta \rangle}^r(\gamma) = \begin{cases} f^q(\gamma) & \text{if } \gamma < \text{ot}(S_\alpha^q) \\ \tau + \delta & \text{if } \gamma = \text{ot}(S_\alpha^q) + \delta, \delta < \beta, \end{cases}$$

and so the same also holds true with r replaced everywhere by p . We therefore have that

$$x \Vdash^p \langle \lambda^p, \text{ot}(S^p \cap \tau) + \beta \rangle \Vdash^p \langle \kappa, \tau + \beta \rangle,$$

and

$$\pi_{x \langle \lambda^p, \text{ot}(S^p \cap \tau) + \beta \rangle}^p = (f^p)^{-1} \circ \pi_{x \langle \kappa, \tau + \beta \rangle}^p.$$

Recalling that $p' \stackrel{\lambda^p}{\sim} p$ and considering $f^{p'}$, we may conclude that

$$x \Vdash^{p'} \langle \lambda^p, \text{ot}(S^p \cap \tau) + \beta \rangle \Vdash^{p'} \langle \kappa, f^q(\text{ot}(S_\alpha^q) + \beta) \rangle = y,$$

with

$$\pi_{xy}^{p'} = f^{p'} \circ (f^p)^{-1} \circ \pi_{x \langle \kappa, \tau + \beta \rangle}^p.$$

If $\gamma < \text{ot}(S_\alpha^q)$, then

$$\pi_{xy}^{p'}(\gamma) = f^{p'} \circ (f^p)^{-1}(f^q(\gamma)) = f^q(\gamma).$$

If $\gamma = \text{ot}(S_\alpha^q) + \delta$, then

$$\begin{aligned} \pi_{xy}^{p'}(\gamma) &= f^{p'} \circ (f^p)^{-1}(\tau + \delta) \\ &= f^{p'}(\text{ot}(S^p \cap \tau) + \delta) \\ &= f^q(\text{ot}(S_\alpha^q) + \delta) \\ &= f^q(\gamma). \end{aligned}$$

Hence, we have $x \Vdash^{p'} y$ with $\pi_{xy}^{p'} = f^q \upharpoonright (o(x) + 1) = \pi_{xy}^q$, and therefore the same is true in s .

In either case $s \leq q$ as claimed, and so r is indeed a reduction of q to $\mathbb{P}_{\kappa, \alpha}$.

Of course there is more to complete embeddings than the existence of reductions. The inclusion $\mathbb{P}_{\kappa, \alpha} \hookrightarrow \mathbb{P}$ certainly preserves the relation \leq ; we must also show that it preserves incompatibility. So suppose p and q in $\mathbb{P}_{\kappa, \alpha}$ are compatible in \mathbb{P} with some common extension $r \in \mathbb{P}$. Let r' be the unique condition such that $\lambda^{r'} = \lambda^r$, $r' \stackrel{\lambda^r}{\sim} r$, and $S^{r'} = S^r \cap \alpha$, as given by Lemma 22. Clearly $r' \in \mathbb{P}_{\kappa, \alpha}$, and it is straightforward to verify that because $p \leq r$ and $q \leq r$, $p \leq r'$ and $q \leq r'$. So the inclusion $\mathbb{P}_{\kappa, \alpha} \hookrightarrow \mathbb{P}$ indeed preserves incompatibility, and hence is a complete embedding.

Finally, suppose that $\alpha > \kappa$ but $\text{cf}(\alpha) < \kappa$. Let $q \in \mathbb{P} \setminus \mathbb{P}_{\kappa, \alpha}$ be such that $S^q \setminus \alpha \neq \emptyset$ and $S^q \cap \alpha$ is cofinal in α (using Lemmas 18 and 27 and Proposition 24 as usual). We claim that q has no reduction to $\mathbb{P}_{\kappa, \alpha}$. Suppose for the sake of contradiction that p is a reduction of q to $\mathbb{P}_{\kappa, \alpha}$, and let r be a common extension of p and q in \mathbb{P} . As in the argument above for incompatibility preservation, let r' be the condition obtained from r but truncating S^r to $S^r \cap \alpha$.

We again obtain $r' \leq p$, so r' is also a reduction of q to \mathbb{P} . Also, if q' is likewise obtained from q by truncating S^q to $S^q \cap \alpha$, then it is straightforward to check that $q' \geq r'$ (indeed, $q' \geq q \geq r$, so the verification for incompatibility preservation also applies to this case). In particular, $S^{r'} \supseteq S^q \cap \alpha$, and for any $\langle \lambda^q, \delta \rangle$ in q with $\delta < \text{ot}(S^q \cap \alpha)$, $\langle \lambda^q, \delta \rangle \Vdash^{r'} \langle \kappa, f^q(\delta) \rangle$.

Let r'' be obtained from r' by applying Lemma 28, so that in particular $\theta_{\lambda^{r''}}^{r''} = \text{ot}(S^{r''})$ and $S^{r''} = S^{r'}$. Since the set of $f^q(\delta)$ for $\delta < \text{ot}(S^q \cap \alpha)$ is cofinal in $\alpha = \sup(S^{r'})$, it follows that the set

$$\{\zeta \mid \exists \delta < \text{ot}(S^q \cap \alpha) (\langle \lambda^q, \delta \rangle \Vdash^{r''} \langle \lambda^{r''}, \zeta \rangle \Vdash^{r''} \langle \kappa, f^q(\delta) \rangle)\}$$

is unbounded in $\theta_{\lambda^{r''}}^{r''}$.

Now suppose that s is a common extension of q and r'' with $\lambda^s > \lambda^{r''}$ (using Lemma 28 again if necessary). Let $\beta = \text{ot}(S^q \cap \alpha)$; then there is some $\tau \geq \alpha$ such that $\langle \lambda^q, \beta \rangle \Vdash^q \langle \kappa, \tau \rangle$, and hence $\langle \lambda^q, \beta \rangle \Vdash^s \langle \kappa, \tau \rangle$. Since $\lambda^{r''}$ is a mangal of s , let γ be such that $\langle \lambda^q, \beta \rangle \Vdash^s \langle \lambda^{r''}, \gamma \rangle \Vdash^s \langle \kappa, \tau \rangle$. We must have $\gamma < \theta_{\lambda^{r''}}^s = \theta_{\lambda^{r''}}^{r''}$, so there is some $\delta < \text{ot}(S^q \cap \alpha)$ and $\zeta > \gamma$ such that $\langle \lambda^q, \delta \rangle \Vdash^{r''} \langle \lambda^{r''}, \zeta \rangle \Vdash^{r''} \langle \kappa, f^q(\delta) \rangle$. Now

$$\pi_{\langle \lambda^{r''}, \gamma \rangle \langle \kappa, \tau \rangle}^s \circ \pi_{\langle \lambda^q, \beta \rangle \langle \lambda^{r''}, \gamma \rangle}^s(\delta) = \pi_{\langle \lambda^q, \beta \rangle \langle \kappa, \tau \rangle}^s(\delta) = \pi_{\langle \lambda^q, \beta \rangle \langle \kappa, \tau \rangle}^q(\delta) = f^q(\delta)$$

So by axiom M.2, $\langle \lambda^{r''}, \pi_{\langle \lambda^q, \beta \rangle \langle \lambda^{r''}, \gamma \rangle}^s(\delta) \rangle \Vdash^s \langle \kappa, f^q(\delta) \rangle$. But $\pi_{\langle \lambda^q, \beta \rangle \langle \lambda^{r''}, \gamma \rangle}^s(\delta) < \gamma < \zeta$, and we already have that $\langle \lambda^{r''}, \zeta \rangle \Vdash^s \langle \kappa, f^q(\delta) \rangle$, giving a contradiction. Thus, there cannot be a reduction p of q to $\mathbb{P}_{\kappa, \alpha}$, and hence the inclusion $\mathbb{P}_{\kappa, \alpha} \hookrightarrow \mathbb{P}$ is not a complete embedding. \dashv

§6. Homogeneity. When we force to obtain a morass in the presence of various large cardinal axioms, we will sometimes be able to preserve the large cardinal property by simply requiring that our generic be below some master condition. Of course, for each cardinal κ with the given large cardinal property, there will be a different master condition. Hence, to be able to preserve the large cardinal property for all of these cardinals at once, one would like to be able to impose each master condition “after the fact”, substituting it into the “bottom” of any given generic to give a new generic suited to the large cardinal at hand. This is achieved with Theorem 46 of this section.

In some forcing posets, such as the standard ones to force \diamond and \square , compatibility implies comparability, and further one of the conditions will be an initial segment of the other. When these initial segments may be freely interchanged giving new conditions, a great deal of homogeneity for the partial order results.

While we do not have these same properties in \mathbb{P} , our notion of μ -equivalence is an appropriate substitute for comparability which does follow from compatibility (Lemma 21), with the part of the morass condition below μ being a suitable “initial segment” for us

to interchange. To be precise, we will use μ -equivalence in defining automorphisms on suitably chosen open dense suborders of \mathbb{P} , with the automorphism being a sort of “initial segment interchange”. We will then show in Theorem 46 that such automorphisms may be used to construct, within an arbitrary generic extension $V[G]$, \mathbb{P} -generics over V containing any given condition. The definition of these automorphisms is the crux of our argument for Theorem 46, and by extension, the large cardinal preservation arguments in Sections 7 and 8. Velleman’s partial order of [12], with no mangal requirement for extension, lacks this crucial homogeneity.

First, we define the suborders of interest.

DEFINITION 37. *For any $\alpha < \kappa$, let*

$$\mathbb{P}^\alpha = \{p \in \mathbb{P} \mid \lambda^p \geq \alpha\}.$$

PROPOSITION 38. *For all $\alpha < \kappa$, \mathbb{P}^α is open dense in \mathbb{P} .*

PROOF. Density follows from Lemma 28 and Proposition 24, and openness is immediate from the definition of \leq . \dashv

Of course, this means that generics for \mathbb{P}^α are easily translatable into generics for \mathbb{P} and conversely, while preserving the extension universe $V[G]$.

We will define our automorphisms by “interchanging” initial pieces of conditions.

DEFINITION 39. *Let p, q and r in \mathbb{P} be three conditions such that $\lambda^q \geq \lambda^p = \lambda^r$, $\theta_{\lambda^p}^p = \theta_{\lambda^r}^r$, and $q \stackrel{\lambda^p}{\sim} p$. Define q_r^p as follows: let*

$$\theta_\alpha^{q_r^p} = \begin{cases} \theta_\alpha^r & \text{if } \alpha \leq \lambda^p \\ \theta_\alpha^q & \text{if } \alpha \geq \lambda^p \end{cases}$$

$$S^{q_r^p} = S^q$$

$$\begin{aligned} x \Vdash^{q_r^p} y \iff & (l(x) < l(y) \leq \lambda^p \wedge x \Vdash^r y) \vee \\ & (\lambda^p \leq l(x) < l(y) \wedge x \Vdash^q y) \vee \\ & (l(x) < \lambda^p < l(y) \wedge \\ & \exists z \in \mathcal{S}^q(l(z) = \lambda^p \wedge x \Vdash^r z \Vdash^q y)), \end{aligned}$$

and for $x \Vdash^{q_r^p} y$,

$$\pi_{xy}^{q_r^p} = \begin{cases} \pi_{xy}^r & \text{if } l(x) < l(y) \leq \lambda^p \\ \pi_{xy}^q & \text{if } \lambda^p \leq l(x) < l(y) \\ \pi_{zy}^q \circ \pi_{xz}^r & \text{if } l(z) = \lambda^p \wedge x \Vdash^r z \Vdash^q y. \end{cases}$$

Intuitively q_r^p is just q with p replaced by r , although S^p and S^r are irrelevant for the construction.

PROPOSITION 40. *For p, q and r as in Definition 39, q_r^p is a morass condition and $q_r^p \stackrel{\lambda^p}{\sim} r$.*

PROOF. It is straightforward to check that all of the requirements for a morass condition hold for q_r^p , because they hold for q and r . In particular, note that M.2 holds for $l(x) < \lambda^p < l(y)$ because $\pi_{xy}^{q_r^p}$ is defined by composition for such x and y . Also, because we only require M.6 and M.7 to hold for immediate \prec -successors in a morass condition, these two axioms are immediate from the fact that they hold in q and r . It is immediate from the definition that λ^p is also a mangal of the new condition q_r^p , and $\lambda^p = \lambda^r$ so it is also a mangal of r . Since the requisite equalities trivially hold, it follows that $q_r^p \stackrel{\lambda^p}{\sim} r$. \dashv

DEFINITION 41. Let p and r in \mathbb{P} be such that $\lambda^p = \lambda^r$ and $\theta_{\lambda^p}^p = \theta_{\lambda^r}^r$. Then define $\varphi^{p,r} : \mathbb{P}^{\lambda^p} \rightarrow \mathbb{P}^{\lambda^p}$ by

$$\varphi^{p,r}(q) = \begin{cases} q_r^p & \text{if } q \stackrel{\lambda^p}{\sim} p \\ q_p^r & \text{if } q \stackrel{\lambda^p}{\sim} r \\ q & \text{otherwise.} \end{cases}$$

Note first that this is well defined: if $p \stackrel{\lambda^p}{\sim} q \stackrel{\lambda^p}{\sim} r$, then $p \stackrel{\lambda^p}{\sim} r$, and it is easy to see that $q_r^p = q_p^r = q$.

Because of the mangal requirement for the relation $\stackrel{\lambda^p}{\sim}$, $\varphi^{p,r}$ is self-inverse: $\varphi^{p,r^2}(q) = q$ for all $q \in \mathbb{P}^{\lambda^p}$. Note that this is the point in the argument which necessitates the mangal requirement, and concomitantly where the definition of \leq in Velleman's partial order fails for our purposes. We could have defined q_r^p without such a requirement, and would still have obtained a morass condition, but the corresponding φ would not have been surjective. The range of such a φ would not include any condition that was "the same as p up to level λ^p " but which did not have λ^p as a mangal, and likewise for r . Since the set of such conditions is not codense in \mathbb{P}^{λ^p} , this is a real cause for concern, hence the introduction of mangals.

PROPOSITION 42. For conditions p and r satisfying the requirements of Definition 41, the function $\varphi^{p,r}$ is an automorphism of the poset \mathbb{P}^{λ^p} .

PROOF. Since $\varphi^{p,r}$ is a self-inverse bijection, it only remains to show that $\varphi^{p,r}$ respects \leq . So suppose that $q \leq s \in \mathbb{P}^{\lambda^p}$. Then $q \stackrel{\lambda^s}{\sim} s$ and $\lambda^p < \lambda^s$, so $q \stackrel{\lambda^p}{\sim} p \leftrightarrow s \stackrel{\lambda^p}{\sim} p$ and $q \stackrel{\lambda^p}{\sim} r \leftrightarrow s \stackrel{\lambda^p}{\sim} r$ by Lemma 20 (note that this would not necessarily be the case if we were to use compatibility in place of the $\stackrel{\lambda^p}{\sim}$ relation).

For the sake of the argument at hand, we may therefore assume without loss of generality that $q \stackrel{\lambda^p}{\sim} p \stackrel{\lambda^p}{\sim} s$, so that $\varphi^{p,r}(q) = q_r^p$ and $\varphi^{p,r}(s) = s_r^p$. To see that λ^q remains a mangal of $\varphi^{p,r}(s)$, note by Proposition 40 that $\lambda^p = \lambda^r$ is a mangal of s_r^p , so any edge of s_r^p factors as an edge in r followed by an edge in s from level λ^p , the latter

factorising through level λ^q since λ^q is a mangal of s . The remainder of the verification that $q_r^p \leq s_r^p$ is trivial from the definitions. \dashv

The following lemma will also be useful.

LEMMA 43. *If p and r are conditions in \mathbb{P} such that $\lambda^p = \lambda^r$ and $\theta_{\lambda^p}^p = \theta_{\lambda^r}^r$, and further $S^p = S^r$, then $\varphi^{p,r}(p) = r$ and $\varphi^{p,r}(r) = p$.*

PROOF. Of course $\varphi^{p,r}$ is symmetric in p and r , so it suffices to consider one case. The set S^r determines f^r and hence also \prec^r and π_{xy}^r up from level λ^r , and $\varphi^{p,r}(p) = p_r^p \stackrel{\lambda^r}{\sim} r$. \dashv

With these tools at our disposal, we are almost ready to prove the main theorem of this section. The idea of the proof of Theorem 46 is to find a condition r in G large enough to “cover” a given p , so that we can use $\varphi^{p,r}$ to “patch in” p in place of r in the generic. The following two lemmas allow us to find a suitable such r .

LEMMA 44. *For any subset X of κ^+ such that $|X| < \kappa$, the set*

$$D_X = \{r \in \mathbb{P} \mid X \subset S^r\}$$

is open dense in \mathbb{P} .

PROOF. Density is obvious by combining Lemma 27 with Proposition 24. Openness is immediate from the definition of extension. \dashv

LEMMA 45. *For any ordinal $\mu < \kappa$, the set*

$$E_\mu = \{r \in \mathbb{P} \mid \exists \alpha > \text{ot}(S^r)(\lambda^r = \mu + \omega^{\text{ot}(S^r)} \cdot \alpha) \wedge \theta_{\lambda^r}^r = \text{ot}(S^r)\}$$

is dense in \mathbb{P} .

PROOF. First note that μ only affects the minimum value of λ for elements of E_μ , and not its “form”: it is a simple result of ordinal arithmetic, that for any μ, γ and α_0 , there is an α_1 such that $\mu + \omega^\gamma \cdot \alpha_0 = \omega^\gamma \cdot \alpha_1$. Now given any p in \mathbb{P} , we may simply use Lemma 28 repeatedly, “gluing together” with Proposition 24 (noting that the argument there will not increase S), to give an extension in E_μ . \dashv

We are now ready to prove the main theorem of this section.

THEOREM 46 (Homogeneity). *Let G be \mathbb{P} -generic over V and let p be a condition in \mathbb{P} . Then there exists a $G' \subset V[G]$ such that $p \in G'$ and G' is also \mathbb{P} -generic over V . Moreover, $V[G'] = V[G]$.*

Note that we write $G' \subset V[G]$ rather than $G' \in V[G]$ because we are not assuming that \mathbb{P} is set sized from the perspective of the ground model V — this will be relevant in Section 8 below.

PROOF. By Lemmas 44 and 45, there is some $r \in G \cap D_{S^p} \cap E_{\lambda^p}$. We claim that p may be extended to a condition p' such that $S^{p'} = S^r$, $\lambda^{p'} = \lambda^r$, and $\theta_{\lambda^{p'}}^{p'} = \theta_{\lambda^r}^r$. Note that in Lemma 27, the extension q of p constructed satisfies $\lambda^q = \lambda^p + \omega^{\text{ot}(S^q)}$. This construction is to add a single new limit ordinal to S , whereas we wish to add the whole of $S^r \setminus S^p$. However, we may construct a sequence of conditions q_α ,

whereby if $\langle \sigma_\alpha \mid \alpha < \beta \rangle$ is the increasing enumeration of limit ordinals in $S^r \setminus S^p$, $q_{\alpha+1}$ is obtained from q_α by the method of Lemma 27 adding σ_α and its finitary successors, $q_0 = p$, and q_λ for limit ordinals λ is obtained by the method of Proposition 24 to close off with a new top level. Using Proposition 24 one last time if β is a limit ordinal and taking $q_{\beta-1}$ otherwise, we get a lower bound q for the sequence such that $S^q = S^r$ and

$$\lambda^q = \lambda^p + \sum_{0 < \alpha < \beta} \omega^{\text{ot}(S^{q_\alpha})}.$$

Now, the summation on the right (excluding λ^p) is clearly less than or equal to $\omega^{\text{ot}(S^q)} \cdot \beta$. Further, β must be less than or equal to $\text{ot}(S^q)$, so we have that

$$\lambda^q \leq \lambda^p + \omega^{\text{ot}(S^q)} \cdot \text{ot}(S^q).$$

But now

$$\lambda^r = \lambda^p + \omega^{\text{ot}(S^r)} \cdot \delta = \lambda^p + \omega^{\text{ot}(S^q)} \cdot \delta \quad \text{for some } \delta > \text{ot}(S^q),$$

so noting as in Lemma 45 that adding $\omega^{\text{ot}(S^q)}$ to an ordinal yields an ordinal divisible by $\omega^{\text{ot}(S^q)}$, we see that we may extend q further to a p' with $S^{p'} = S^r$ and $\lambda^{p'} = \lambda^r$, by simply applying Lemma 28 and Proposition 24 repeatedly. Also, since we must use Lemma 28 at least once to get p' , we will further have that $\theta_{\lambda^{p'}}^{p'} = \text{ot}(S^{p'}) = \text{ot}(S^r) = \theta_{\lambda^r}^r$, and so p' has all of the desired attributes in common with r .

Now in $V[G]$, let G^{λ^r} denote $G \cap \mathbb{P}^{\lambda^r}$, let $G_{p'}^{\lambda^r}$ denote $\varphi^{p',r} \text{``} G^{\lambda^r}$, and let

$$G' = \{q \in \mathbb{P} \mid \exists s \in G_{p'}^{\lambda^r} (s \leq q)\}.$$

Because $\varphi^{p',r}$ is a poset automorphism and \mathbb{P}^{λ^r} is open dense in \mathbb{P} , G' is \mathbb{P} -generic over V , and $V[G'] = V[G]$. Because $\varphi^{p',r}(r) = p'$ and $r \in G^{\lambda^r}$, $p' \in G_{p'}^{\lambda^r}$, and so $p \in G'$, as desired. \dashv

§7. Preserving large cardinals. In [7], it was shown that using a reverse Easton iteration of forcings to construct morasses, one may force morasses to exist at every cardinal while preserving a hyperstrong or n -superstrong cardinal, for $0 < n \leq \omega$. However, except in the case of 1-superstrong cardinals, this result is only achieved for a single cardinal, as the generic used must fall below a specific master condition that is dependent on the embedding witnessing n -superstrength or hyperstrength, and so a single generic cannot be used to preserve the large cardinal strength of many different cardinals. On the other hand, with our Homogeneity Theorem (Theorem 46), we can circumvent this problem, expressing for each master condition p the extension $V[G]$ as $V[G']$ for some $G' \ni p$. We give the details of the argument in this section.

Assume the GCH holds (this can be forced while preserving large cardinals as in [7] and Section 2).

DEFINITION 47. *The Global Mangrove Partial Order R is the reverse Easton iteration of partial orders \dot{Q}_α , where \dot{Q}_α names \mathbb{P}_α if α is a regular uncountable cardinal, and \dot{Q}_α names the trivial forcing otherwise.*

Note that \mathbb{P}_κ will continue to denote the forcing to add a mangrove at κ , whereas R_κ will now denote the iteration up to stage κ of such forcings.

THEOREM 48. *If $V \models ZFC + GCH$ and G is R -generic over V , then*

$$V[G] \models ZFC + GCH +$$

“There is a mangrove at κ for every regular cardinal κ .”

PROOF. For each regular cardinal κ , we have that \mathbb{P}_κ is κ -closed and κ^+ -cc (Propositions 24 and 26), and moreover $|\mathbb{P}_\kappa| = (\kappa^+)^{\kappa} = \kappa^+$. As mentioned in Subsection 4.4, this implies that forcing with \mathbb{P}_κ preserves cardinals, and a nice names argument gives that the GCH is preserved as well.

Applying the Factor Lemma, we may factorise R as $R_\kappa * \dot{R}^\kappa$ for any κ , with R^κ forced to be κ -closed. As discussed in the proof of Lemma 2, this implies that R is tame.

We prove that cardinals and the GCH are preserved by induction on the iteration length. Successor stages are immediate from the observations about \mathbb{P}_κ above.

At singular limit cardinal λ stages, we observe by factorising R_λ as $R_{\kappa^+} * R^{\kappa^+}$ that each cardinal $\kappa < \lambda$ is preserved, whence λ itself is also preserved, and moreover the GCH is preserved below λ . The iteration R_λ has a dense suborder of size $\prod_{\kappa < \lambda} \kappa^+ = \lambda^+$, so cardinals greater than λ^+ are preserved, and the GCH holds at and above $(\lambda^+)^V$. Finally, taking unions over $\kappa < \lambda$ of nice names for subset of κ , we obtain λ^+ nice names for subsets of λ , so there is a surjection from $(\lambda^+)^V$ to $(2^\lambda)^{V[G_\lambda]}$. Hence, λ^+ is also preserved, as is the GCH at κ .

At regular limit stages λ , where we take direct limits, the argument is similar but easier, since R_λ will have a dense suborder of size $\sum_{\kappa < \lambda} \kappa^+ = \lambda$. Indeed if λ is Mahlo, we even have that R_λ satisfies the λ chain condition — see [1], Corollary 2.4. In any case, we have shown that every stage of the iteration preserves cardinals and the GCH, and so since the closedness of R^κ increases with κ , R itself preserves cardinals and the GCH.

Showing that mangroves exist at every regular uncountable cardinal of the generic extension is now straightforward. Given a regular uncountable κ and factorising R as $R_\kappa * \dot{\mathbb{P}}_\kappa * \dot{R}^{\kappa+1}$, we have that there is a mangrove at κ after the first $\kappa + 1$ stages of the iteration. Now the statement “ M is a mangrove at κ ” is absolute for models containing M , κ as a regular cardinal, and κ^+ as its successor. Thus,

since the final part $R^{\kappa+1}$ of the iteration is κ^+ -closed, the mangrove at κ will remain a mangrove after this final part of the iteration, and we are done. \dashv

THEOREM 49. *Suppose that $V \models ZFC + GCH$, and G is R -generic over V . Then for any $n \in \omega + 1$ and any cardinal κ , if*

$$V \models \kappa \text{ is } n\text{-superstrong}$$

then

$$V[G] \models \kappa \text{ is } n\text{-superstrong}.$$

Similarly, if κ is hyperstrong in V , then κ is hyperstrong in $V[G]$.

PROOF. As mentioned above, the case of 1-superstrongs is proven in [7]. The proofs in the other cases also follow exactly as in that paper, with the simple modification that because of our homogeneity theorem, a generic $G' \subset V[G]$ with $V[G'] = V[G]$ may be chosen containing any given master condition. Now, we should be careful at this point, because for n -superstrength with $n > 1$, we will want to take a master condition at not just a single iterand of the forcing, but many. Factorise R as

$$R_\kappa * \dot{\mathbb{P}}_\kappa * R^{(\kappa, j(\kappa))} * \dots * \mathbb{P}_{j^n(\kappa)} * R^{j^n(\kappa)+1},$$

and let G be R -generic over V . We wish to modify (within $V[G]$) the upper part of G so as to contain the master conditions for the lower part. There are no restrictions on $G_{j(\kappa)}$. We can modify G so that $G(j(\kappa))$ contains the master condition from $G(\kappa)$. Now $R^{(j(\kappa), j^2(\kappa))}$ is $j(\kappa)^+$ -directed-closed since each of its iterands is (see [1], Theorem 2.7), so we may take a single master condition in $R^{(j(\kappa), j^2(\kappa))}$ determined by $G^{(\kappa, j(\kappa))}$. By arguments as in [5], our iteration $R^{(j(\kappa), j^2(\kappa))}$ is sufficiently homogeneous to allow us to modify $G^{(j(\kappa), j^2(\kappa))}$, and hence G , to lie below this single master condition. Repeating all of this another $n - 2$ times, we construct a $G'_{j^{n-1}(\kappa)} \subset V[G]$ below all of the necessary master conditions to make the arguments of [7] go through. Of course, if $n = \omega$ the question arises as to whether this is legitimate, but since an indirect limit is taken at stage $j^\omega(\kappa)$ the methods of [5] are still applicable.

Therefore, each n -superstrong or hyperstrong cardinal is preserved in *every* generic extension by R , and so any generic extension $V[G]$ preserves all such cardinals. \dashv

§8. 1-extendible cardinals and morasses. In the case of forcing the GCH in Section 2, there was no major difficulty in proving that 1-extendible cardinals were preserved. In particular, the fact that any embedding j witnessing 1-extendibility lifted presented no great problems, essentially because the forcing affecting the domain of j was small — P_κ was a set in H_{κ^+} . To show that 1-extendible cardinals are preserved while forcing morasses to exist, we will have

to work significantly harder, as the forcing of interest, $R_{\kappa+1}$, will be class sized from the point of view of H_{κ^+} .

THEOREM 50. *Let V be a model of ZFC. There is class generic extension $V[G]$ of V satisfying the GCH, having mangroves at every regular cardinal in $V[G]$, and such that every 1-extendible cardinal of V is 1-extendible in $V[G]$.*

PROOF. We of course apply the two stage iteration $P * R$ of the GCH Partial Order from Section 2 followed by the Global Mangrove Partial Order above; recall that whilst taking infinite iterations of class forcings is generally problematic, there is no difficulty with 2-stage iterations. We have already seen in Theorems 6 and 48 that if $G_g * G_m$ is $P * R$ -generic over V , then $V[G_g * G_m]$ will satisfy ZFC + GCH and have mangroves at every regular cardinal. It therefore only remains to show that any 1-extendible cardinal κ of V will remain 1-extendible in $V[G_g * G_m]$.

So suppose κ is 1-extendible in V , and let $j : H_{\kappa^+}^V \rightarrow H_{\lambda^+}^V$ be an elementary embedding witnessing the 1-extendibility of κ . Let j^* be the lift of j to $H_{\kappa^+}^{V[G_g]}$, as obtained in Theorem 6, so j^* witnesses that κ is 1-extendible in $V[G_g]$. By Theorem 9 we may assume that G_g is such that j^* is elementary from $\langle H_{\kappa^+}^{V[G_g]}, G_g(\kappa) \rangle$ to $\langle H_{\lambda^+}^{V[G_g]}, G_g(\lambda) \rangle$ as \mathcal{L}_{STG} -structures, without changing $V[G_g]$.

Now the Global Mangrove Partial Order R may be factorised as $R \cong R_{\kappa+1} * \dot{R}^{\kappa+1}$, with $\dot{R}^{\kappa+1}$ forced to be κ^+ -closed. Hence,

$$H_{\kappa^+}^{V[G_g * G_m]} = H_{\kappa^+}^{V[G_g * G_m, \kappa+1]},$$

where $G_{m, \kappa+1} = G_m \upharpoonright R_{\kappa+1}$.

To lift our embedding

$$j^* : H_{\kappa^+}^{V[G_g]} \rightarrow H_{\lambda^+}^{V[G_g]}$$

to

$$j^{**} : H_{\kappa^+}^{V[G_g * G_m]} \rightarrow H_{\lambda^+}^{V[G_g * G_m]},$$

we will first lift it to $H_{\kappa^+}^{V[G_g * G_m, \kappa]}$, and then again to $H_{\kappa^+}^{V[G_g * G_m, \kappa+1]} = H_{\kappa^+}^{V[G_g * G_m]}$.

Lifting j^* to $j^{*'} : H_{\kappa^+}^{V[G_g * G_m, \kappa]} \rightarrow H_{\lambda^+}^{V[G_g * G_m, \kappa]}$ is like the case of forcing the GCH while preserving 1-extendible cardinals (that is, Theorem 6) and at first glance seems to pose no problems. In particular, $H_{\kappa^+}^{V[G_g * G_m, \kappa]} = H_{\kappa^+}^{V[G_g]}[G_m, \kappa]$, and $j^* \upharpoonright R_\kappa$ is the identity function, so the Lifting Lemma gives us that $j^{*'} : H_{\kappa^+}^{V[G_g]}[G_m, \kappa] \rightarrow H_{\lambda^+}^{V[G_g]}[G_m, \lambda]$ is elementary. However, we actually want to demonstrate that $j^{*'}$ is elementary from

$$\langle H_{\kappa^+}^{V[G_g]}[G_m, \kappa], G_g(\kappa), G_m, \kappa \rangle$$

to

$$\langle H_{\lambda^+}^{V[G_g]}[G_m, \lambda], G_g(\lambda), G_m, \lambda \rangle.$$

The Lifting Lemma will certainly accommodate this if the forcing relation is definable for pre-existing class predicates A (thus dealing with $G_g(\kappa)$; $G_{m,\kappa}$ is not a problem since R_κ is still only set sized from the perspective of $H_{\kappa^+}^{V[G_g]}$). But we must be careful, as the definition of $\Vdash^* A$ as given for example in [6], page 34, relies on our ground model having a definable stratification into sets.

There are (at least) two ways to deal with this. First, we could redefine $p \Vdash^* A(\sigma)$, analogously to the definition of $p \Vdash^* G(\sigma)$ in the proof of Lemma 8. Specifically, it is not hard to check that

$$p \Vdash^* A(\sigma) \iff \forall q \leq p \exists r \leq q \exists a \in A(r \Vdash^* \sigma = \check{a})$$

is an appropriate definition.

Alternatively, we can give away the punch line of the proof that the next lift goes through, and observe that our ground model $\langle H_{\kappa^+}^{V[G_g]}, G_g(\kappa) \rangle$ *does* have a definable stratification into sets. Specifically, every element of $H_{\kappa^+}^{V[G_g]} = H_{\kappa^+}^{V[G_g, \kappa]}$ is coded into the Cohen generic $G_g(\kappa)$, and so $H_{\kappa^+}^{V[G_g]} = L_{\kappa^+}[G_g(\kappa)]$. Hence, the $L_\alpha[G_g(\kappa)]$ hierarchy is a stratification of $H_{\kappa^+}^{V[G_g]}$ into sets, $p \Vdash A(\sigma)$ is definable, and the Lifting Lemma gives that $j^{*'}$ is indeed elementary from $\langle H_{\kappa^+}^{V[G_g]}[G_{m,\kappa}], G_g(\kappa), G_{m,\kappa} \rangle$ to $\langle H_{\lambda^+}^{V[G_g]}[G_{m,\lambda}], G_g(\lambda), G_{m,\lambda} \rangle$. That is, we have

$$j^{*'} : \langle H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}, G_g(\kappa), G_{m,\kappa} \rangle \rightarrow \langle H_{\lambda^+}^{V[G_g * G_{m,\lambda}]}, G_g(\lambda), G_{m,\lambda} \rangle$$

elementary.

We next wish to lift $j^{*'}$ to $j^{**} : H_{\kappa^+}^{V[G_g * G_{m,\kappa+1}]} \rightarrow H_{\lambda^+}^{V[G_g * G_{m,\lambda+1}]}$. Since \mathbb{P}_κ enjoys the κ^+ -cc, every \mathbb{P}_κ -name in $V[G_g * G_{m,\kappa}]$ has an equivalent name in $H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}$, so once again this is the same as lifting to $H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}[G_m(\kappa)]$. The forcing partial order \mathbb{P}_κ is class-sized from the point of view of $H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}$, so we need a stratification of $H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}$ into sets in $H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}$ in order to deduce from Theorem 2.18 of [6] that the forcing relation is definable, and subsequently apply the Lifting Lemma. Note on the other hand that the pretameness of \mathbb{P}_κ is unproblematic for us: we know (from the V perspective) that forcing with \mathbb{P}_κ over $H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}$ will give us a model of ZFC^- , $H_{\kappa^+}^{V[G_g * G_{m,\kappa+1}]}$, so since pretameness is implied by ZF^- preservation (Proposition 2.17 of [6]), \mathbb{P}_κ is pretame. Alternatively, observe that the κ^+ -cc directly implies that \mathbb{P}_κ is pretame.

Now we have $H_{\kappa^+}^{V[G_g]} = L_{\kappa^+}[G_g(\kappa)]$, so

$$H_{\kappa^+}^{V[G_g * G_{m,\kappa}]} = H_{\kappa^+}^{V[G_g]}[G_{m,\kappa}] = L_{\kappa^+}[G_g(\kappa), G_{m,\kappa}].$$

Thus, the $L_\alpha[G_g(\kappa), G_{m,\kappa}]$ hierarchy is a stratification of $H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}$ into sets, which moreover is definable in $\langle H_{\kappa^+}^{V[G_g * G_{m,\kappa}]}, G_g(\kappa), G_{m,\kappa} \rangle$. So we can conclude that the forcing relation is definable, and by the Lifting Lemma, if

$$j^{*'} \text{``} G_m(\kappa) \subseteq G_m(\lambda),$$

then j^{**} lifts to an elementary

$$j^{**} : H_{\kappa^+}^{V[G_g * G_m, \kappa]}[G_m(\kappa)] \rightarrow H_{\lambda^+}^{V[G_g * G_m, \lambda]}[G_m(\lambda)]$$

as desired. But now $|G_m(\kappa)| = \kappa^+$ and $G_m(\kappa)$ is directed, so since \mathbb{P}_λ is λ -directed-closed (Lemma 24), there is a lower bound $p \in \mathbb{P}_\lambda$ for $j^{**} \restriction G_m(\kappa)$. By Theorem 46, there is a $G'_m(\lambda) \subset H_{\lambda^+}^{V[G_g * G_m, \lambda]}[G_m(\lambda)]$ which is \mathbb{P}_λ -generic over $H_{\lambda^+}^{V[G_g * G_m, \lambda]}$, contains the master condition p , and such that

$$H_{\lambda^+}^{V[G_g * G_m, \lambda]}[G_m(\lambda)] = H_{\lambda^+}^{V[G_g * G_m, \lambda]}[G'_m(\lambda)].$$

Therefore, without changing the model $H_{\lambda^+}^{V[G_g * G_m, \lambda+1]}$ we may apply the Lifting Lemma, and obtain an elementary embedding

$$j^{**} : H_{\kappa^+}^{V[G_g * G_m, \kappa+1]} \rightarrow H_{\lambda^+}^{V[G_g * G_m, \lambda+1]}.$$

As noted above, $H_{\kappa^+}^{V[G_g * G_m, \kappa+1]} = H_{\kappa^+}^{V[G_g * G_m]}$, and likewise with κ replaced by λ , so j^{**} witnesses that κ is 1-extendible in $V[G_g * G_m]$. \dashv

§9. Universal morasses. We now consider a notion extending that of a $(\kappa, 1)$ -morass, in which a predicate at the top of the morass, built up through the morass, codes every subset of κ . These are of interest to practitioners of pcf theory for use in the construction of scales.

DEFINITION 51. *An augmented κ -morass is a $(\kappa, 1)$ -morass M , along with a set A_x associated to each element x of \mathcal{S}^M , such that*

1. *for every $x \in \mathcal{S}^M$, $A_x \subseteq o(x)$,*
2. *for $w, x \in \mathcal{S}^M$ with $l(w) = l(x)$ and $o(w) < o(x)$, $A_w = A_x \cap o(w)$, and*
3. *for $x, y \in \mathcal{S}^M$ with $x \prec^M y$, $A_x = (\pi_{xy}^M)^{-1} \restriction A_y$.*

For any augmented morass $\langle M, A \rangle$, we denote by A_κ the set $\bigcup_{\tau < \kappa^+} A_{\langle \kappa, \tau \rangle} \subseteq \kappa$. A κ -universal morass is an augmented κ -morass $\langle M, A \rangle$ such that $\mathcal{P}(\kappa) \subset L[A_\kappa]$.

It turns out that by modifying the mangrove forcing appropriately, we may force a κ -universal morass. The idea is this: assuming we already have a subset A'_κ of κ^+ such that $\mathcal{P}(\kappa) \subset L[A'_\kappa]$, we may force an augmented morass with A'_κ at the top level. However, there are new subsets of κ added by forcing the universal morass itself. Thus, we also want the morass to be self-coding, giving a subset of κ^+ at the top level which encodes the morass itself.

For those familiar with Jensen's definition of a morass, as in [4] for example, this will seem entirely straightforward: at each node of the morass we can encode the morass below that node, and then the morass maps will naturally preserve this information, cohering at level κ of the morass to give a subset of κ^+ encoding the morass itself. However, in the definition of a morass that we have used

(Definition 11, taken from [12]), there is no requirement that the morass maps are this “nice”. We are confident that by finding the right balance between Jensen’s definition of a morass and Velleman’s, this elegant solution might be made to go through. This will have the added benefit that attempts to make our suborder homogeneous will have one less problem to worry about: since the augmentation of the morass for self-coding can be thought of as occurring after the forcing takes place, it will not interfere with any desired homogeneity. On the other hand, new ideas will in any case be necessary to make our suborder homogeneous while coding up all of the ground model subsets of κ .

In the meantime, we use a different technique to resolve the matter of self-coding. In essence, we shall force our morass to be self-coding by treating the final code for the morass much like the pre-existing subset A'_κ . To do this, we place the restriction on our conditions that they are only allowed to hit morass coding points at which membership or not is decided by the condition itself. The top level subset of the augmented morass that we force will code the given A'_κ at successor ordinals, and the morass itself will be coded at 0 and limit ordinals, so that there will be no problems with S^p being closed under ordinal successors and predecessors for conditions p . For terminological convenience, we shall consider 0 to be a limit ordinal for the rest of this article. Note that the function $\omega \cdot : \alpha \mapsto \omega \cdot \alpha$ is the increasing enumeration of all limit ordinals.

In defining our coding we shall work a little harder than is necessary for our main goal, in order to show how close one can get to homogeneity for our partial order, and where it can nevertheless fail. For this, we wish to avoid coding redundant information. It turns out that mangals will be helpful for this, and moreover the following particular kind.

DEFINITION 52. *Given a morass or morass condition M , a determined mangal of M is a mangal β of M such that*

1. θ_β is a limit ordinal, and
2. for all $y \in M$ with $l(y) = \beta$, there is a $\tau < \kappa^+$ such that $y \Vdash \langle \kappa, \tau \rangle$.

We will need the following basic result stemming from the morass axioms, showing that no “splitting” occurs at limit stages of the tree.

LEMMA 53. *Let M be a morass or morass condition, and let $\beta \leq \kappa$ or λ^M respectively. Then for all $y \neq y' \in M$ with $l(y) = l(y') = \beta$ and $o(y) + 1 < \theta_\beta$ and $o(y') + 1 < \theta_\beta$, there are $x \neq x' \in M$ such that $x \Vdash y$, $x' \Vdash y'$, and x and x' are \Vdash -incomparable.*

PROOF. Take y and y' with $l(y) = l(y') = \beta$ and $o(y') < o(y)$. By M.5, there is an $x \Vdash y$ and a $\nu < o(x)$ such that $\pi_{xy}(\nu) = o(y')$, whence from M.2 we have that $\langle l(x), \nu \rangle \Vdash y'$. Taking $x' = \langle l(x), \nu \rangle \Vdash y'$, then, we are done. \dashv

LEMMA 54. *Suppose β is a determined mangal of M . Then the data θ_β , the tree structure given by \prec up to level β and directly from level β to level κ , and the corresponding maps π , are all determined by the part of the morass below level β and the morass maps from such lower nodes to level κ .*

PROOF. First we reconstruct θ_β . Let $X \subseteq \kappa^+$ be the set of $\tau \in \kappa^+$ such that there is an $x \in M$ with $l(x) < \beta$ and $x \prec \langle \kappa, \tau \rangle$, and define an equivalence relation \sim on X by

$$\sigma \sim \tau \leftrightarrow \forall x (l(x) < \beta \rightarrow (x \prec \langle \kappa, \sigma \rangle \leftrightarrow x \prec \langle \kappa, \tau \rangle)).$$

Then by Lemma 53, θ_β is in bijection with the set of \sim -equivalence classes, and moreover thanks to M.2, θ_β is the order type of the set of \sim -equivalence classes ordered by least element. For any $\gamma < \theta_\beta$ and $\tau \in \kappa^+$, $\langle \beta, \gamma \rangle \prec \langle \kappa, \tau \rangle$ if and only if τ is in the γ -th \sim -equivalence class of X , and similarly for any $x \in M$ with $l(x) < \beta$, $x \prec \langle \beta, \gamma \rangle$ if and only if there is a $\tau < \kappa^+$ in the γ -th equivalence class of X such that $x \prec \langle \kappa, \tau \rangle$.

We also claim that the corresponding maps π may be recovered. For $x \prec \langle \beta, \gamma \rangle$ this is clear: if $\mu < o(x)$ and $\nu < \gamma$, then $\pi_{x \langle \beta, \gamma \rangle}(\mu) = \nu$ if and only if for $\tau < \kappa^+$ such that $x \prec \langle \kappa, \tau \rangle$, $\pi_{x \langle \kappa, \tau \rangle}(\mu)$ is in the ν -th equivalence class of X . The maps $\pi_{\langle \beta, \gamma \rangle \langle \kappa, \tau \rangle}$ for $\langle \beta, \gamma \rangle \prec \langle \kappa, \tau \rangle$ may now be deduced from the Commutativity morass axiom: for any $\nu < \gamma$ we have by M.5 that there is an $x \prec \langle \beta, \gamma \rangle$ and a $\mu < o(x)$ such that $\pi_{x \langle \beta, \gamma \rangle}(\mu) = \nu$, and then $\pi_{\langle \beta, \gamma \rangle \langle \kappa, \tau \rangle}(\nu)$ must be $\pi_{x \langle \kappa, \tau \rangle}(\mu)$. \dashv

Because of this “reconstructibility” property, when we code up the morass we need only flag the determined mangals as such, rather than all of the corresponding details. Thus, when we iteratively construct conditions that are sufficiently closed with respect to hitting the morass coding points, we will be able to add on the necessary new top level without adding condition-specific morass coding point information.

We are now ready to define our encoding for the morass. The first $\kappa \cdot \kappa$ morass coding points will be used to encode the values θ_α for $\alpha < \kappa$. The next $\kappa \cdot \kappa$ will be used to encode the morass relation \prec below level κ , and the next κ^4 will be used to encode the maps π_{xy} below level κ . The rest of the morass coding points will be used to encode the relation $x \prec y$ and maps π_{xy} for y with level κ .

DEFINITION 55. *Let p be a morass condition, and for all $\gamma \leq \lambda^p + 1$ define*

$$\begin{aligned} ND(\gamma) &= \{\alpha < \gamma \mid \alpha \text{ is not a determined mangal of } p\} \\ \iota(\gamma) &= \sum_{\alpha < \gamma} \theta_\alpha, \quad \text{and} \\ \delta(\gamma) &= \sum_{\alpha \in ND(\gamma)} \theta_\alpha. \end{aligned}$$

Let $m : \iota(\lambda^p + 1) \rightarrow \mathcal{S}^p$ $n : \delta(\lambda^p + 1) \rightarrow \mathcal{S}^p$ be the enumeration in lexicographic order of the nodes of p with level in $\lambda^p + 1$ and $ND(\lambda^p + 1)$ respectively. Let $b : \kappa^+ \times \kappa^+ \rightarrow \kappa^+$ be the standard bijection according to the max-lex order on $\kappa^+ \times \kappa^+$. Then the morass code for p is a partial function

$$c^p : \{\omega \cdot \alpha \mid \alpha \in \kappa^+\} \rightarrow 2$$

such that the following hold.

1. For all $\alpha \in ND(\lambda^p + 1)$ and $\zeta < \kappa$,

$$c^p(\omega \cdot (\kappa \cdot \alpha + \zeta)) = \begin{cases} 1 & \text{if } \zeta < \theta_\alpha \\ 0 & \text{otherwise.} \end{cases}$$

For every β a determined mangal of p and $\zeta < \kappa$, $c^p(\omega \cdot (\kappa \cdot \beta + \zeta)) = 0$.

2. For $\alpha < \iota(\lambda^p)$, let μ be the least determined mangal of p greater than $l(m(\alpha))$ if such exists, and $\lambda^p + 1$ otherwise. Then for all $\zeta < \iota(\mu)$,

$$c^p(\omega \cdot (\kappa^2 + \kappa \cdot \alpha + \zeta)) = \begin{cases} 1 & \text{if } m(\alpha) \Vdash^p m(\zeta) \\ 0 & \text{otherwise.} \end{cases}$$

If μ is a determined mangal, then for all $\zeta \geq \iota(\mu)$, $c^p(\omega \cdot (\kappa^2 + \kappa \cdot \alpha + \zeta)) = 0$.

3. For all $\alpha < \iota(\lambda^p)$, all $\zeta < \iota(\mu)$ with μ as in item 2, and all $\eta < \kappa$,

$$\begin{aligned} & c^p(\omega \cdot (\kappa^2 \cdot 2 + \kappa^2 \cdot \alpha + \kappa \cdot \zeta + \eta)) \\ &= \begin{cases} 1 & \text{if } m(\alpha) \Vdash^p m(\zeta) \text{ and } \eta \in \pi_{m(\alpha)m(\zeta)} " (o(m(\alpha)) + 1) \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

If μ is a determined mangal, then for all $\zeta \geq \iota(\mu)$ and $\eta < \kappa$,

$$c^p(\omega \cdot (\kappa^2 \cdot 2 + \kappa^2 \cdot \alpha + \kappa \cdot \zeta + \eta)) = 0.$$

4. For all $\tau \in \mathcal{S}^p$, $\alpha < \delta(\lambda^p + 1)$, and $\nu < \kappa^+$,

$$\begin{aligned} & c^p(\omega \cdot (\kappa^3 + \kappa \cdot b((\tau, \nu)) + \alpha)) \\ &= \begin{cases} 1 & \text{if } n(\alpha) \Vdash^p \langle \kappa, \tau \rangle \text{ and } \nu \in \pi_{n(\alpha)\langle \kappa, \tau \rangle} " (o(n(\alpha)) + 1) \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

5. At all other limit ordinals c^p is undefined.

Clearly the entire condition p may be reconstructed from the information provided by c^p . Also, our coding scheme is nice with respect to the mangrove \leq relation:

LEMMA 56. If p and q are morass conditions with $p \leq q$ in the mangrove forcing, then $c^q \subseteq c^p$.

PROOF. If new nodes, tree edges and attendant maps are added to extend q to p , these will be coded at limit ordinals on which c^q was left undefined. If new tree edges from nodes in q to nodes at level κ are added, then the node at level κ must be new (by injectivity of f^p) and again the code is only extended at points where it was previously undefined. \dashv

Now we must force in a way that respects c^p and our predicate A'_κ , and moreover such that we will be able to respect c^q for extensions q . The modification to the quagmire forcing necessary to achieve this is given by the following definition. As usual, for any subset X of κ^+ we denote by χ_X the characteristic function of X .

DEFINITION 57. *Let A' be a subset of κ^+ such that $\mathcal{P}(\kappa) \subset L[A']$, and let $\chi_{A'}$ be its characteristic function. Let p be a morass condition. Let $\chi_A^p : \kappa^+ \rightarrow \kappa^+$ be the partial function defined by*

$$\chi_A^p(\tau) = \begin{cases} \chi_{A'}(\nu) & \text{if } \tau = \nu + 1 \\ c^p(\tau) & \text{if } c^p \text{ is defined on } \tau \end{cases}$$

(and undefined otherwise), and let A_κ^p denote the subset of $\text{dom}(\chi_A^p)$ with characteristic function χ_A^p . The condition p is a universal morass condition for A' , or simply universal morass condition when A' is clear from the context, if following properties hold.

1. *The set $S^p \cap \text{Lim}$, where Lim denotes the class of limit ordinals, is a subset of $\text{dom}(c^p)$.*
2. *For all $x \in S^p$, if τ and ν in S^p are such that $x \Vdash^p \langle \kappa, \tau \rangle$ and $x \Vdash^p \langle \kappa, \nu \rangle$, then $(\pi_{x(\kappa, \tau)}^p)^{-1} \langle A_\kappa^p \cap \tau \rangle = (\pi_{x(\kappa, \nu)}^p)^{-1} \langle A_\kappa^p \cap \nu \rangle$.*
3. *For all $x \in S^p$ with $l(x) \leq \lambda^p$, there is a $y \in p$ such that $l(y) = \kappa$ and $x \Vdash^p y$.*

Let U denote the set of all universal morass conditions for A' . The Universal Morass Partial Order \mathbb{U}_κ for A' is the suborder of \mathbb{P}_κ consisting of all universal morass conditions for A' , that is, $\mathbb{U}_\kappa = \langle U, \leq \rangle$ where \leq is defined as for the mangrove forcing.

As with \mathbb{P} , we will generally omit the subscript κ from \mathbb{U}_κ .

We claim that forcing with \mathbb{U} yields a universal morass. The corresponding set A_κ will be $\bigcup_{p \in G} A_\kappa^p$, and the other sets $A(x)$ for x in the generic morass will “filter down” from this top level predicate: if x, y are nodes in the generic morass with $l(y) = \kappa$, then $A(x) = \pi_{xy}^{-1} \langle A_\kappa \cap o(y) \rangle$. Requirements 2 and 3 ensure that $A(x)$ will be well-defined for all x in the generic morass, and the coherence properties necessary to have an augmented morass will be automatic.

The verification that forcing with \mathbb{U} still yields a morass is much as in Section 4, with a little extra work to check that requirements 1–3 of Definition 57 do not cause problems. We shall now walk through the modified versions of the enunciations of Section 4, only describing the necessary modifications to the old proofs. In particular, the fact

that straightforward verifications go through will not be mentioned.

Lemma 18'. *There is a universal morass condition q with $S^q = \omega$.*

PROOF. The construction is exactly as for Lemma 18. Since $\langle 0, 0 \rangle$ must be in q , $c^q(0)$ is defined (and equal to 1). Every node of the q constructed is below a unique node at level κ , and so q satisfies all of the requirements of Definition 57. \dashv

In Subsection 4.3 regarding μ -equivalence, Lemmas 20 and 21 go through unchanged, but Lemma 22 now fails in general. However, we do not require that lemma for our present purposes.

Proposition 24' *The poset \mathbb{U} is κ -directed-closed.*

PROOF. As in the proof of Proposition 24, given a directed set Y of conditions, we take the union of the conditions and add a new top level if necessary. The requirement of Definition 57 hold below the new top level because they do for every member of Y , and 2 and 3 hold at the new top level by its definition. \dashv

We need to suitably strengthen the hypotheses of Lemma 25 to make the proof go through in the context of \mathbb{U} .

Lemma 25' *Let p and q be universal morass conditions such that $\lambda^p = \lambda^q$ and $p \stackrel{\lambda^p}{\sim} q$. Suppose further that there is some $S^0 \subset \kappa^+$ such that both S^p and S^q end-extend S^0 , $\min(S^q \setminus S^0) \geq \sup(S^p)$, and $(f^p)^{-1} \restriction A_\kappa^p = (f^q)^{-1} \restriction A_\kappa^q$. Then p and q are compatible in \mathbb{U} .*

PROOF. The added hypothesis ensures that requirement 2 holds of the condition constructed in the proof of Lemma 25, and the other two requirements are immediate. \dashv

Proposition 26' *The poset \mathbb{U} is κ^+ -cc.*

PROOF. At the end of the Δ -system argument for Proposition 26, we may add one extra stage of thinning, to ensure that $(f^p)^{-1} \restriction A_\kappa^p$ is the same for all of the conditions p in the set in question, thus making Lemma 25' applicable. \dashv

It will be convenient to consider the variants of Lemmas 27 and 28 in reverse order.

Lemma 28' *For any $p \neq \mathbf{1} \in \mathbb{U}$ there is a $q \leq p$ in \mathbb{U} such that $S^q = S^p$, $\theta_{\lambda^q}^q = \text{ot}(S^q)$, and $\lambda^q = \lambda^p + \omega^{\text{ot}(S^q)}$.*

PROOF. It is immediate that the q constructed in the proof of Lemma 28 is a universal morass condition. \dashv

Lemma 27' *For any $p \neq \mathbf{1} \in \mathbb{U}$ and any limit ordinal $\sigma < \kappa^+$, there is a $q \leq p$ in \mathbb{U} such that $S^q \supseteq ((\sigma + \omega) \setminus \sigma)$.*

PROOF. The only point of concern is to ensure that $\sigma \in \text{dom}(c^q)$. If $\sigma < \kappa^3$, we may first apply Lemma 28' repeatedly, "gluing together" with Proposition 24', to obtain a p' with sufficiently large $\lambda^{p'}$ and $\delta(\lambda^{p'})$ that $c^{p'}$ is defined on σ . The desired extension q may then be constructed starting from p' as in the proof of Lemma 27.

If $\sigma = \omega \cdot (\kappa^3 + \kappa \cdot b((\tau, \nu)) + \alpha)$ for some τ, ν and α , then $\tau \leq \sigma$. We may again make a preliminary extension of p to a p' with

$\alpha < \delta(\lambda^{p'} + 1)$. If $\tau = \sigma$, then the q constructed in the proof of Lemma 27 satisfies $\sigma \in \text{dom}(c^q)$, and is thus a universal morass condition. Otherwise, by induction we may extend p' to p'' with $\tau \in S^{p''}$, which implies that $\sigma \in c^{p''}$, and so constructing q from this starting point again makes it a universal morass condition. \dashv

We now have all the necessary ingredients for the following analogue of Theorem 29, which we believe merits its own number.

THEOREM 58. *Forcing with \mathbb{U}_κ yields a κ -universal morass.*

PROOF. Let G be \mathbb{U} -generic over V . The morass part M of the universal morass will simply be the direct limit of the conditions in G , as was the case in our original mangrove forcing. The proof that M is indeed a morass proceeds exactly as for Theorem 29.

As discussed earlier, we set $A_\kappa = \bigcup_{p \in G} A_\kappa^p$, and define $A(x)$ for every $x \in M$ by

$$A(x) = \begin{cases} A_\kappa \cap o(x) & \text{if } l(x) = \kappa \\ \pi_{xy}^{-1} " A(y) & \text{if } x \prec y \text{ and } l(y) = \kappa. \end{cases}$$

By requirements 2 and 3 this uniquely defines $A(x) \subseteq o(x)$ for every $x \in M$. Requirement 2 for an augmented morass is satisfied thanks to morass axiom M.2 and our definition of $A(x)$, and requirement 3 for an augmented morass is immediate from the Commutativity morass axiom.

Therefore, to prove that $\langle M, A \rangle$ so defined is a κ -universal morass in $V[G]$, it only remains to show that every subset of κ in $V[G]$ lies in $L[A_\kappa]$. We have by assumption that $\mathcal{P}(\kappa)^V \subset L[A']$, and A' is coded into A_κ on successor ordinals. Now since \mathbb{U} has the κ^+ chain condition, every element of $\mathcal{P}(\kappa)^{V[G]}$ has a nice name in $H_{\kappa^+}^V$, and therefore in $L[A_\kappa]$. But now the morass M is entirely encoded into A_κ on limit ordinals, and the generic G may readily be reconstructed from the morass M as the set of all universal morass conditions that are substructures of M . Hence, $G \subset L[A_\kappa]$, so within $L[A_\kappa]$ we can correctly evaluate names, and therefore $\mathcal{P}(\kappa)^{V[G]} \subset L[A_\kappa]$. Overall then, we have that $\langle M, A \rangle$ indeed constitutes a universal morass in $V[G]$. \dashv

We may of course also take a reverse Easton iteration of these partial orders \mathbb{U}_κ to force a κ -universal morass to exist at every uncountable regular cardinal κ . For this, we will need to interleave Cohen forcings to obtain the sets A' with the forcing posets \mathbb{U}_κ .

DEFINITION 59. *The Global Universal Morass Partial Order W is the reverse Easton iteration forcing with $\text{Fn}(\kappa^+, 2, \kappa^+) * \mathbb{U}_\kappa$ at uncountable regular cardinal stages κ , and the trivial forcing elsewhere. In the former case, the A' from which the partial order \mathbb{U}_κ is defined is taken to be the subset of κ^+ whose characteristic function is the union of the generic for the preceding $\text{Fn}(\kappa^+, 2, \kappa^+)$.*

THEOREM 60. *Forcing with the Global Universal Morass Partial Order yields a universe in which the GCH holds and universal morasses exist at every uncountable regular cardinal.*

PROOF. This is a straightforward verification as for Theorem 48. Also note that we only need the GCH to hold below κ^+ for the lemmas leading up to Theorem 58, so we need not even start with a model of the GCH. \dashv

Now we consider large cardinal preservation.

THEOREM 61. *Forcing with the Global Universal Morass Partial Order W preserves all superstrong cardinals of the ground model. Moreover, given any 1-extendible, hyperstrong or n -superstrong cardinal κ of the ground model for $n \in \omega + 1$, the W -generic G may be chosen so as to preserve the large cardinal strength of κ .*

PROOF. For n -superstrong and hyperstrong cardinals this is exactly as for Theorem 12 of [7]. For 1-extendible cardinals, the situation is much as in Theorem 50. Given $j : H_{\kappa^+} \rightarrow H_{\lambda^+}$, we first lift to $j' : H_{\kappa^+}^{V[G_\kappa]} \rightarrow H_{\lambda^+}^{V[G_\lambda]}$ using the fact that W_κ is a set forcing from the perspective of H_{κ^+} . Then we lift to

$$j'' : \langle H_{\kappa^+}^{V[G_\kappa * G_{g,\kappa}]}, G_{g,\kappa} \rangle \rightarrow \langle H_{\lambda^+}^{V[G_\lambda * G_{g,\lambda}]}, G_{g,\lambda} \rangle,$$

where $G_{g,\kappa}$ and $G_{g,\lambda}$ are the generics for $\text{Fn}(\kappa^+, 2, \kappa^+)$ and $\text{Fn}(\lambda^+, 2, \lambda^+)$ respectively, arguing as in the proof of Theorem 9. Finally, since \mathbb{U}_λ is λ -directed-closed, we may choose the generic G so that the corresponding part $G_{U,\lambda}$ lies below $j''G_{U,\kappa}$, and hence such that our embedding lifts to

$$j^* : H_{\kappa^+}^{V[G_\kappa * G_{g,\kappa} * G_{U,\kappa}]} = H_{\kappa^+}^{V[G]} \rightarrow H_{\lambda^+}^{V[G_\lambda * G_{g,\lambda} * G_{U,\lambda}]} = H_{\lambda^+}^{V[G]},$$

as required. \dashv

As a final point, we consider briefly the question of the homogeneity, or lack thereof, of \mathbb{U} . With our carefully defined coding scheme, we can find conditions that seem to almost be sufficiently interchangeable to define an automorphism of \mathbb{U} as was done in Definition 41 for \mathbb{P} . Let $\text{supp}(c)$ denote the subset of $\text{dom}(c)$ on which c is not 0.

PROPOSITION 62. *Let p and q be universal morass conditions. Then there are universal morass conditions $p' \leq p$ and $q' \leq q$ such that $\lambda^{p'} = \lambda^{q'}$, $\theta_{\lambda^{p'}}^{p'} = \theta_{\lambda^{q'}}^{q'}$, $\text{dom}(c^{p'}) = \text{dom}(c^{q'})$, $\lambda^{p'}$ is a determined mangal of both p' and q' , and $\text{supp}(c^{p'}) \subseteq S^{p'} = S^{q'} \supseteq \text{supp}(c^{q'})$.*

PROOF. Having put in the effort to set up our coding carefully in Definition 55, this is now relatively easy. Note that the fact that $\theta_{\lambda^{p'}}^{p'} = \theta_{\lambda^{q'}}^{q'}$ follows from $S^{p'} = S^{q'}$ and the requirement that every node of a universal morass condition lie below a node at level κ .

Now, we may construct sequences p_i and q_i for $i < \omega$ with $p_0 = p$, $q_0 = q$, and for every $i \in \omega$,

$$\text{supp}(c^{p_i}) \cup \text{supp}(c^{q_i}) \subseteq S^{p_{i+1}} \cap S^{q_{i+1}}$$

and

$$\text{dom}(c^{p_i}) \cup \text{dom}(c^{q_i}) \subseteq \text{dom}(c^{p_{i+1}}) \cap \text{dom}(c^{q_{i+1}}).$$

Note that this latter requirement implies that $\lambda^{p_{i+1}} \geq \max(\lambda^{p_i}, \lambda^{q_i}) \leq \lambda^{q_{i+1}}$. The construction is by combining Lemmas 27' and 28' and Proposition 24'; note that in the proofs of all three of these results, the condition constructed has a determined mangrove as the top level, so in particular parts 2 and 3 of the definition of the morass code (Definition 55) do not pose a problem. Applying Proposition 24' to obtain lower bounds for the sequences $(p_i)_{i \in \omega}$ and $(q_i)_{i \in \omega}$ gives the desired p' and q' respectively, since the new top level in each case is a determined mangal, requiring no new 1s in the morass code. \dashv

Unfortunately, this is still not sufficient to be able to define an automorphism interchanging p' and q' . The difficulty lies in the fact that an extension r of p' may be able to “see” extra information about the structure of p' through the required agreement between $(\pi_{xy}^r)^{-1} “A_\kappa^r$ and $(\pi_{xz}^r)^{-1} “A_\kappa^r$ whenever $x \prec^r y$ and $x \prec^r z$ with $l(y) = l(z) = \kappa$. Of course, this may contradict what “happens in” q , thus blocking the existence of an automorphism of \mathbb{U} interchanging p' and q' . So we do not have the requisite homogeneity needed to make Theorem 61 go through for a proper class of n -superstrong or hyperstrong cardinals, and must leave this as an open question. However, as mentioned above, it seems likely that a solution to this problem will emerge with a suitable compromise between the Jensen and Velleman definitions of a morass.

REFERENCES

- [1] JAMES E. BAUMGARTNER, *Iterated forcing*, **Surveys in set theory**, London Mathematical Society Lecture Note Series, no. 87, Cambridge University Press, 1983, pp. 1–59.
- [2] JAMES CUMMINGS, *Iterated forcing and elementary embeddings*, To appear in *The Handbook of Set Theory*.
- [3] JAMES CUMMINGS and ERNEST SCHIMMERLING, *Indexed squares*, **Israel Journal of Mathematics**, vol. 131 (2002), pp. 61–99.
- [4] KEITH J. DEVLIN, **Constructibility**, Springer, Berlin, 1984.
- [5] NATASHA DOBRINEN and SY D. FRIEDMAN, *Homogeneous iteration and measure one covering relative to HOD*, To appear.
- [6] SY D. FRIEDMAN, **Fine structure and class forcing**, de Gruyter Series in Logic and Its Applications, no. 3, de Gruyter, Berlin, 2000.
- [7] ———, *Large cardinals and L-like universes*, **Set theory: Recent trends and applications** (Alessandro Andretta, editor), Quaderni di Matematica, vol. 17, Seconda Università di Napoli, 2005, pp. 93–110.
- [8] THOMAS JECH, **Set theory**, Third Millenium ed., Springer, 2003.
- [9] AKIHIRO KANAMORI, **The higher infinite**, 2nd ed., Springer, 2003.
- [10] KENNETH KUNEN, **Set theory**, North-Holland, 1980.
- [11] LEE STANLEY, *Review of papers by Dan Velleman*, **The Journal of Symbolic Logic**, vol. 54 (1989), no. 2, pp. 639–646.

[12] DANIEL J. VELLEMAN, *Morasses, diamond, and forcing*, *Annals of Mathematical Logic*, vol. 23 (1982), no. 2-3, pp. 199–281.

[13] ———, *Simplified morasses*, *The Journal of Symbolic Logic*, vol. 49 (1984), no. 1, pp. 257–271.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF BRISTOL
UNIVERSITY WALK
BRISTOL, BS8 1TW, UK
E-mail: Andrew.Brooke-Taylor@bristol.ac.uk

KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC
UNIVERSITY OF VIENNA
WÄHRINGER STRASSE 25
A-1090, WIEN, AUSTRIA
E-mail: sdf@logic.univie.ac.at